arXiv:0911.2646v2 [math-ph] 13 Jan 2010

On Universality of Bulk Local Regime of the Deformed Laguerre Ensemble

T. Shcherbyna
Institute for Low Temperature Physics, Kharkov, Ukraine.
E-mail: t_shcherbina@rambler.ru

2000 Mathematics Subject Classification. Primary 15A52; Secondary 15A57

Abstract

We consider the deformed Laguerre Ensemble $H_n = \dfrac{1}{m}\Sigma_n^{1/2}A_{m,n}A_{m,n}^*\Sigma_n^{1/2}$ in which Σ_n is a positive hermitian matrix (possibly random) and $A_{m,n}$ is a $n\times m$ complex Gaussian random matrix (independent of Σ_n), $\dfrac{m}{n}\to c>1$. Assuming that the Normalized Counting Measure of Σ_n converges weakly (in probability) to a non-random measure $N^{(0)}$ with a bounded support we prove the universality of the local eigenvalue statistics in the bulk of the limiting spectrum of H_n.

1 Introduction.

Universality is an important topic of the random matrix theory. It deals with statistical properties of eigenvalues of $n\times n$ random matrices on intervals whose length tends to zero as $n\to\infty$. According to the universality conjecture these properties do not depend to large extent on the ensemble. The conjecture was proposed by Dyson in the early 60s. To formulate it we need some notations and definitions. Denote by $\lambda_1^{(n)},\dots,\lambda_n^{(n)}$ the eigenvalues of the random matrix. Define the normalized eigenvalue counting measure (NCM) of the matrix as

$$N_n(\triangle)=\sharp\{\lambda_j^{(n)}\in\triangle, j=\overline{1,n}\}/n,\quad N_n(\mathbb{R})=1, \qquad (1.1) \qquad \boxed{\texttt{NCM}}$$

where $\triangle$ is an arbitrary interval of the real axis. For many known random matrices the expectation $\overline{N}_n=\mathbf{E}\{N_n\}$ is absolutely continuous, i.e.,

$$\overline{N}_n(\triangle)=\int\limits_{\triangle}\rho_n(\lambda)d\,\lambda. \qquad (1.2) \qquad \boxed{\texttt{rhon}}$$

The non-negative function ρ_n in (1.2) is called the density of states.

Define also the k-point correlation function $R_k^{(n)}$ by the equality:

$$\mathbf{E}\left\{\sum_{j_1\neq\ldots\neq j_k}\varphi_k(\lambda_{j_1},\dots,\lambda_{j_k})\right\}=\int\varphi_k(\lambda_1,\dots,\lambda_k)R_k^{(n)}(\lambda_1,\dots,\lambda_k)d\lambda_1,\dots,d\lambda_k, \qquad (1.3) \qquad \boxed{\texttt{R}}$$

where $\varphi_k:\mathbb{R}^k\to\mathbb{C}$ is bounded, continuous and symmetric in its arguments and the summation is over all k-tuples of distinct integers $j_1,\dots,j_k\in\{1,\dots,n\}$. Here and below integrals without limits denote the integration over the whole real axis.

The behavior of N_n as $n \to \infty$ is studied for many ensembles. It is shown that N_n converges weakly to a non-random limiting measure N. The limiting measure is normalized to unity and as a rule is absolutely continuous

$$N(\mathbb{R}) = 1, \quad N(\triangle) = \int\limits_{\triangle} \rho(\lambda) d\,\lambda. \tag{1.4}$$

The non-negative function ρ in (1.4) is called the limiting density of states of the ensemble.

We will call the spectrum the support of N and define the bulk of the spectrum as

$$\operatorname{bulk} N = \{\lambda | \exists (a,b) \subset \operatorname{supp} N : \lambda \in (a,b), \inf_{\mu \in (a,b)} \rho(\mu) > 0\}.$$

Then the universality hypothesis on the bulk of the spectrum says that for $\lambda_0 \in \operatorname{bulk} N$ we have:

(i) for any fixed k uniformly in $\xi_1, \xi_2, \dots, \xi_k$ varying in any compact set in $\mathbb{R}$

$$\lim_{n\to\infty} \frac{1}{(n\rho_n(\lambda_0))^k} R_k^{(n)} \left(\lambda_0 + \frac{\xi_1}{\rho_n(\lambda_0)\, n}, \dots, \lambda_0 + \frac{\xi_k}{\rho_n(\lambda_0)\, n} \right) = \det\{S(\xi_i - \xi_j)\}_{i,j=1}^k, \tag{1.5}$$

where

$$S(x) = \frac{\sin \pi x}{\pi x}, \tag{1.6}$$

and $R_k^{(n)}$ and ρ_n are defined in (1.3) and (1.2);

(ii) if

$$E_n(\triangle) = \mathbf{P}\{\lambda_i^{(n)} \notin \triangle, \, i = \overline{1,n}\}, \tag{1.7}$$

is the gap probability, then

$$\lim_{n\to\infty} E_n \left(\left[\lambda_0 + \frac{a}{\rho_n(\lambda_0)\, n}, \lambda_0 + \frac{b}{\rho_n(\lambda_0)\, n} \right] \right) = \det\{1 - S_{a,b}\}, \tag{1.8}$$

where the operator $S_{a,b}$ is defined on $L_2[a,b]$ by the formula

$$S_{a,b} f(x) = \int\limits_a^b S(x-y) f(y) d\,y,$$

and S is defined in (1.6).

Bulk universality was proved initially for ensembles with Gaussian entries (see [11]). Then at the end of 90's it was proved for unitary invariant ensembles of random matrices (known also as unitary matrix models)([1, 2, 3]), end then for Wigner ensemble with some special distribution (see [17]). Recently in the series of important papers [4]-[6] it was shown that local eigenvalue statistics for matrices with independent entries depends only on the first few moments of the distribution of entries. Hence, to prove the universality conjecture it suffices to prove it for the corresponding models with the same moments of the distribution of entries (e.g. for gaussian entries). These results solved the universality problem for Wigner ensemble with subexponential decay of entries.

In this paper we consider the deformed Laguerre Ensemble, i.e. $n \times n$ matrices

$$H_n = \frac{1}{m} \Sigma_n^{1/2} A_{m,n} A_{m,n}^* \Sigma_n^{1/2}, \tag{1.9}$$

where Σ_n is a positive Hermitian $n\times n$ matrix with eigenvalues $\{t_j^{(n)}\}_{j=1}^n \subset \mathbb{R}$ and $A_{m,n}$ is a $n\times m$ matrix, whose entries $\Re a_{\alpha j}$ and $\Im a_{\alpha j}$ are independent Gaussian random variables such that

$$\mathbf{E}\{\Re a_{\alpha j}\} = \mathbf{E}\{\Im a_{\alpha j}\} = 0, \quad \mathbf{E}\{\Re^2 a_{\alpha j}\} = \mathbf{E}\{\Im^2 a_{\alpha j}\} = \frac{1}{2}, \quad \alpha = \overline{1,n},\, j = \overline{1,m}, \tag{1.10}$$ $\boxed{\texttt{A}}$

moreover

$$c_{m,n} := \frac{m}{n} \to c, \quad m,n\to\infty. \tag{1.11}$$ $\boxed{\texttt{c}}$

Let

$$N_n^{(0)}(\triangle) = \sharp\{t_j^{(n)} \in \triangle, j = \overline{1,n}\}/n.$$

be the Normalized Counting Measure of eigenvalues of Σ_n.

The behavior of NCM for the ensemble (1.9) – (1.10) is studied well enough. Indeed, it is easy to see that the spectral properties of matrices (1.9) and of the $m\times m$ matrices $m^{-1}A_{m,n}^*\Sigma_n A_{m,n}$ are closely related. For example, if N_n and N_m^* are the Normalized Counting Measures of (1.1) of these ensembles, then we have

$$N_m^* = (1 - c_{m,n}^{-1})\delta_0 + c_{m,n}^{-1} N_n, \tag{1.12}$$ $\boxed{\texttt{sv}}$

where δ_0 is the Dirac δ-function. For matrices $m^{-1}A_{m,n}^*\Sigma_n A_{m,n}$ it was shown in [7] that if $N_n^{(0)}$ converges weakly with probability 1 to a non-random measure $N^{(0)}$ as $n\to\infty$, then N_m^* also converges weakly with probability 1 to a non-random measure N^*. Moreover the Stieltjes transforms f^* of N^* satisfies the equation

$$f^*(z) = -\left(z - c^{-1}\int \frac{tN^{(0)}(d\,t)}{1 + tf(z)}\right)^{-1}.$$

Hence, we have that if $N_n^{(0)}$ converges weakly to a non-random measure $N^{(0)}$ as $n\to\infty$, then N_n for matrices (1.9) also converges weakly to a non-random measure N, and since according to (1.12) we have that

$$f^*(z) = \frac{1 - c^{-1}}{z} + c^{-1}f(z),$$

where f is the Stieltjes transforms of N, we obtain the equation

$$c^{-1}f(z) + \frac{1 - c^{-1}}{z} = -\left(z - c^{-1}\int \frac{tN^{(0)}(d\,t)}{1 + t(c^{-1}f(z) + (1 - c^{-1})/z)}\right)^{-1}. \tag{1.13}$$ $\boxed{\texttt{eqv\_f}}$

The result on the local regime for the models of the type (1.9) are much more pure. In the case $\Sigma_n = 1$ the universality of the local regime (in the bulk and near the edges of the spectrum) was studied in [8]. The bulk universality for the case $\Sigma_n = 1$, $m/n \to 1$, but $A_{m,n} = A_{m,n}^{(0)} + A_{m,n}^{(1)}$, with gaussian $A_{m,n}^{(0)}$ and $A_{m,n}^{(1)}$ with i.i.d., but not necessary gaussian entries, was studied in [16]. In [15] the limiting distribution of the largest eigenvalue for ensemble (1.9) with $\Sigma_n = I + P$, where P is a finite rank operator, was investigated.

In the present paper we prove universality of the local bulk regime for random matrices (1.9) for a rather general class of Σ_n. The main result is the following theorem.

$\boxed{\texttt{thm:1}}$ **Theorem 1.** *Let $c < 1$ and the eigenvalues $\{t_j^{(n)}\}_{j=1}^n \subset (\mathbb{R}^+)^n$ of Σ_n in (1.9) be a collection of random variables independent of $A_{m,n}$ of (1.10). Assume that there exists a non-random measure $N^{(0)}$ of a bounded support $\sigma \in \mathbb{R}^+$ such that for any finite interval $\Delta \subset \mathbb{R}$ and for any $\varepsilon > 0$*

$$\lim_{n\to\infty} \mathbf{P}_n^{(\tau)}\{|N^{(0)}(\Delta) - N_n^{(0)}(\Delta)| > \varepsilon\} = 0. \tag{1.14}$$ $\boxed{\texttt{conpN0}}$

Then for any $\lambda_0 \in \operatorname{bulk} N$ the universality properties (1.5) and (1.8) hold.

The paper is organized as follows. In Section 2 we give a proof of determinant formulas for correlation functions (1.3) by the method from those of [12, 13]. This formula coincide with formula for the kernel $K_{M,N}$ in [15, Prop.2.1], where the limiting distribution of the largest eigenvalue for ensemble (1.9) for $\Sigma_n = I + P$ with finite rank perturbation P was investigated. Theorem 1 will be proved in Section 3. The method which is used for the limiting transition for the kernel is similar to that of [10]. Section 4 deals with the proof of auxiliary statements for Theorem 1.

Note that we denote by C, C_1, etc. various constants appearing below, which can be different in different formulas.

2 The determinant formulas.

It is well known (see, e.g., [11]) that the correlation functions (1.3) for the GUE $n\times n$ matrix can be written in the determinant form

$$R_k^{(n)}(\lambda_1,\dots,\lambda_k) = \det\{K_n(\lambda_i,\lambda_j)\}_{i,j=1}^k \tag{2.1}$$

with

$$K_n(\lambda_i,\lambda_j) = \sum_{s=0}^{n-1}\phi_s(\lambda_i)\phi_s(\lambda_j), \quad \phi_s(x) = n^{1/4}h_s(\sqrt{n}x)e^{-nx^2/4},$$

where $\{h_s\}_{s\ge 0}$ are orthonormal Hermite polynomials. We want to find analogs of these formulas in the case of random matrices (1.9). To do this we will use the method from [12, 13], where the determinant formulas for deformed Gaussian Unitary Ensemble were derived.

p:ker

Proposition 1. *Let H_n be the random matrix defined in (1.9) and $R_k^{(n)}$ be the correlation function (1.3). Then we have*

$$R_k^{(n)}(\lambda_1,\dots,\lambda_k) = \mathbf{E}_n^{(\tau)}\{\det\{K_n(\lambda_i,\lambda_j)\}_{i,j=1}^k\}, \tag{2.2}$$

Det

with

$$K_n(\lambda,\mu) = -\frac{m}{4\pi^2}\oint_L\oint_\omega \frac{\exp\{m(\lambda u - \mu t)\}\, t^m}{(u-t)u^m}\prod_{j=1}^n\left(\frac{u-\tau_j}{t-\tau_j}\right)dt\,du, \tag{2.3}$$

K

where $\tau_j = 1/t_j^{(n)}$, L is a closed contour, encircling $\{\tau_j\}_{j=1}^n$ and ω is any closed contour encircling 0 and not intersecting L.

The symbol $\mathbf{E}_n^{(\tau)}\{\dots\}$ here and below denotes the expectation with respect to the measure generated by Σ_n.

Proof. The probability distribution $P_n(H_n)$ for ensemble (1.9) is given by (see,e.g.,[15])

$$P_n(H_n) = \frac{1}{Z_n}e^{-m\,\mathrm{tr}\,(\Sigma_n^{-1}H_n)}\det{}^{m-n}H_n, \tag{2.4}$$

p_h

where

$$Z_n = \int e^{-m\,\mathrm{tr}\,(\Sigma_n^{-1}H_n)}\det{}^{m-n}H_n d\,H_n. \tag{2.5}$$

Z

Let us first calculate Z_n. Set

$$\Sigma_n^{-1} = V_0^* L V_0, \quad H_n = V^* X V \tag{2.6}$$

change

where $L = \mathrm{diag}\,(\tau_1, \dots, \tau_n)$, $\tau_j = 1/t_j^{(n)}$, $X = \mathrm{diag}\,(x_1, \dots, x_n)$ and V_0, V are the matrices diagonalizing Σ_n^{-1} and H_n correspondingly. Then the differential $d\,H_n$ in (2.5) transforms to $\triangle^2(X) d\,X d\,\mu(V)$, where $d\,X = \prod\limits_{j=1}^{n} d\,x_j$,

$$\triangle(X) = \prod_{i<j}^{n} (x_i - x_j) \qquad (2.7)$$

VdM

is a Vandermonde determinant, and $\mu(V)$ is the normalized to unity Haar measure on the unitary group $U(n)$. Integral over the unitary group $U(n)$ can be easily computed using the well-known Harish Chandra/Itsykson-Zuber formula (see [11], Appendix 5)

p:Its-Z **Proposition 2.** *Let A and B be normal $n \times n$ matrices with eigenvalues $\{a_i\}_{i=1}^n$, $\{b_i\}_{i=1}^n$, correspondingly. Then we have*

$$\int \exp\{trAU^*BU\} d\,\mu(U) = \frac{\det[\exp\{a_i b_j\}]_{i,j=1}^n}{\triangle(A)\triangle(B)}, \qquad (2.8)$$

Its-Zub

where $\triangle(A)$ and $\triangle(B)$ are Vandermonde determinants for the eigenvalues of A and B.

Shifting $VV_0^* \to V$ and using (2.8) we have from (2.5)

$$Z_n = \int\limits_0^\infty \frac{\det\{e^{-m\tau_j x_k}\}_{j,k=1}^n \triangle(X)}{\triangle(L)} \prod_{j=1}^n x_j^{m-n} d\,X. \qquad (2.9)$$

Since the function under the integral here is symmetric of $\{x_j\}_{j=1}^n$ and $m > n$, we obtain

$$Z_n = \frac{n!}{\triangle(L)} \int\limits_0^\infty e^{-m\sum\limits_{j=1}^n \tau_j x_j} \triangle(X) \prod_{j=1}^n x_j^{m-n} d\,X$$
$$= \frac{n!}{\triangle(L)} \det\left\{ \int\limits_0^\infty e^{-m\sum\limits_{j=1}^n \tau_j x_j} x_j^{m-n+l-1} d\,x_j \right\}_{j,l=1}^n = \frac{n! m^{n(n-1)/2} \prod\limits_{l=1}^n (m-n+l-1)!}{\prod\limits_{j=1}^n (m\tau_j)^m}. \qquad (2.10)$$

Z_n

Let us consider the function

$$U_k(t_1, \dots, t_k) = \mathbf{E}\{\mathrm{tr}\, e^{it_1 H_n} \dots \mathrm{tr}\, e^{it_k H_n}\}.$$

Substituting expression (2.4) we obtain

$$U_k(t_1, \dots, t_k) = \frac{1}{Z_n} \int e^{-m\mathrm{tr}\,(\Sigma_n^{-1} H_n)} \det{}^{m-n} H_n \mathrm{tr}\, e^{it_1 H_n} \dots \mathrm{tr}\, e^{it_k H_n}\, d\,t_1 \dots d\,t_k. \qquad (2.11)$$

U

According to the definition of $R_k^{(n)}$ we have

$$R_k^{(n)}(\lambda_1, \dots, \lambda_k) = \frac{1}{Z_n} \int d\,t_1 \dots d\,t_k\, e^{-i\lambda_1 t_1 - \dots - i\lambda_k t_k} U_k(t_1, \dots, t_k).$$

Doing the change (2.6) and using (2.8) we get

$$\begin{aligned} R_k^{(n)}(\lambda_1, \dots, \lambda_k) \;\; &= \;\; \frac{1}{Z_n \Delta(L)} \prod_{s=1}^k \left(\int e^{-i\lambda_s t_s} d\,t_s \right) \prod_{j=1}^n \left(\int\limits_0^\infty x_j^{m-n} d\,x_j \right) \\ &\quad \times \det\left\{ e^{-mx_j\tau_l} \right\}_{j,l=1}^n \Delta(X) \prod_{s=1}^k \left(\sum_{j=1}^n e^{it_s x_j} \right). \end{aligned} \qquad (2.12)$$

Since the integral in (2.12) is symmetric function of $\{x_l\}_{l=1}^n$, we can write

$$\begin{aligned} R_k^{(n)}(\lambda_1,\dots,\lambda_k) &= \frac{n!}{Z_n\triangle(L)}\sum_{\alpha_1,\dots,\alpha_k=1}^{n}\prod_{s=1}^{k}\Big(\int d\,t_s\Big) \\ &\prod_{j=1}^{n}\left(\int_0^\infty x_j^{m-n}d\,x_j\right) e^{-m\sum\limits_{j=1}^{n}x_j\tau_j+\sum\limits_{s=1}^{k}it_s(x_{\alpha_s}-\lambda_s)}\Delta(X) \end{aligned}$$

It is easy to see, that if some of α_j's coincide (for example, $\alpha_1=\alpha_2=\ldots=\alpha_l$), then the integral over $t_1,\dots,t_l$ in such term becomes $\delta(x_{\alpha_1}-\lambda_1)\prod\limits_{i=1}^{l-1}\delta(\lambda_i-\lambda_{i+1})$ and hence can be omitted for $\lambda_i\neq\lambda_j$. Therefore, integrating over t_s for $\lambda_i\neq\lambda_j$ we obtain

$$\begin{aligned} R_k^{(n)}(\lambda_1,\dots,\lambda_k) &= \frac{n!}{Z_n\triangle(L)}\widetilde{\sum}\prod_{j\notin I_\alpha}\left(\int_0^\infty x_j^{m-n}d\,x_j\right)\prod_{s=1}^{k}\lambda_s^{m-n} \\ &\times\ e^{-m\sum\limits_{j\notin I_\alpha}x_j\tau_j-m\sum\limits_{s=1}^{k}\lambda_s\tau_{\alpha_s}}\triangle(X_\alpha), \end{aligned}$$

where

$$\triangle(X_\alpha)=\triangle(X)\Big|_{x_{\alpha_j}=\lambda_j},\quad j=1,\dots,k$$

and the sum is over all collection $I_\alpha:=\{\alpha_i\}_{i=1}^k$ such that $\alpha_i\neq\alpha_j$. Using $\triangle(X)=\det[x_j^l]_{j,l=0}^{n-1}$ and rewriting this integral as a determinant and computing integrals over x_j we get

$$R_k^{(n)}(\lambda_1,\dots,\lambda_k)=\frac{n!}{Z_n\triangle(L)}\widetilde{\sum}e^{-m\sum\limits_{s=1}^{k}\lambda_s\tau_{\alpha_s}}\det\left\{p_{j,l}^{\alpha}\right\}_{j,l=1}^{n}, \tag{2.13}$$ R1

where

$$p_{j,l}^{\alpha}=\begin{cases}\lambda_s^{m-n+l-1}, & j=\alpha_s \text{ for some } s,\\ \dfrac{(m-n+l-1)!}{(m\tau_j)^{m-n+l}}, & \text{otherwise.}\end{cases} \tag{2.14}$$ p_jl

Note that

$$\lambda_s^{m-n+l-1}=\frac{(m-n+l-1)!m}{2\pi i}\oint\limits_{\omega}\frac{e^{mu\lambda_s}d\,u}{(mu)^{m-n+l}}$$

where ω is a closed contour encircling zero. Hence, (2.13) can be rewritten as

$$\begin{aligned} R_k^{(n)}(\lambda_1,\dots,\lambda_k) &= \frac{n!m^k\prod\limits_{l=1}^{n}(m-n+l-1)!}{Z_n}\widetilde{\sum}\left(\prod_{s=0}^{k}\oint\frac{d\,u_s}{2\pi i}\right) \\ &\times\ \frac{e^{-m\sum\limits_{s=1}^{k}\lambda_s\tau_{\alpha_s}+m\sum\limits_{s=1}^{k}u_s\lambda_s}}{\prod\limits_{j\notin I_\alpha}(m\tau_j)^m\prod\limits_{s=1}^{k}(mu_s)^m}\frac{\det\left\{q_{j,l}^{\alpha}\right\}_{j,l=1}^{n}}{\triangle(L)}, \end{aligned} \tag{2.15}$$

where

$$q_{j,l}^{\alpha}=\begin{cases}(mu_s)^{n-l}, & j=\alpha_s \text{ for some } s,\\ (m\tau_j)^{n-l}, & \text{otherwise.}\end{cases} \tag{2.16}$$ q_jl

It is easy to see that

$$\det\left\{q^{\alpha}_{j,l}\right\}^{n}_{j,l=1}=m^{n(n-1)/2}\prod_{s<t}(u_s-u_t)\prod_{j<l,j,l\notin I_\alpha}(\tau_j-\tau_l)\prod_{s=1}^{k}\prod_{j\notin I_\alpha}\varepsilon_{j,s}(u_s-\tau_j),$$

where

$$\varepsilon_{j,s}=\begin{cases}+1, & \alpha_s<j,\\ -1, & \alpha_s>j.\end{cases}$$

Using this we obtain

$$\frac{\det\left\{q^{\alpha}_{j,l}\right\}^{n}_{j,l=1}}{m^{n(n-1)/2}\triangle(L)}=\frac{\prod\limits_{\alpha_s<\alpha_t}(u_s-u_t)}{\prod\limits_{\alpha_s<\alpha_p}(\tau_{\alpha_s}-\tau_{\alpha_p})}\prod_{s=1}^{k}\prod_{j\notin I_\alpha}\frac{u_s-\tau_j}{\tau_{\alpha_s}-\tau_j}. \tag{2.17}$$ $\boxed{\texttt{frac\_det}}$

Thus, according to the residue theorem (2.15) and (2.10) yield

$$\begin{aligned}R^{(n)}_k(\lambda_1,\dots,\lambda_k)\;&=\;m^k\prod_{s=1}^{k}\oint\limits_{L}\frac{d\,t_s}{2\pi i}\oint\limits_{\omega}\prod_{s=1}^{k}\frac{d\,u_s}{2\pi i}\frac{e^{m\sum\limits_{s=1}^{k}u_s\lambda_s-m\sum\limits_{s=1}^{k}\lambda_s t_s}\prod\limits_{s=1}^{k}t_s^m}{\prod\limits_{s=1}^{k}u_s^m}\\&\times(-1)^{\frac{k(k-1)}{2}}\frac{\prod\limits_{s<t}(u_s-u_t)\prod\limits_{s<p}(t_s-t_p)}{\prod\limits_{s=1}^{k}\prod\limits_{p=1}^{k}(u_s-t_p)}\prod_{s=1}^{k}\prod_{j=1}^{n}\frac{u_s-\tau_j}{t_s-\tau_j},\end{aligned}\tag{2.18}$$

where $\tau_j=1/t^{(n)}_j$, L is a closed contours encircling $\{\tau_j\}^n_{j=1}$ and ω is any closed contour encircling 0 and not intersecting L. Now the identity (see [21], Problem 7.3)

$$(-1)^{\frac{k(k-1)}{2}}\frac{\prod\limits_{s<t}(u_s-u_t)\prod\limits_{s<p}(t_s-t_p)}{\prod\limits_{s=1}^{k}\prod\limits_{p=1}^{s}(u_s-t_p)}=\det\left[\frac{1}{u_s-t_p}\right]^{k}_{p,s=1},$$

and (2.18) yield (2.2) with (2.3). □

3 Proof of Theorem 1.

In this section we prove the universality conjecture (1.5) and (1.8) of the local bulk regime of Hermitian random matrices (1.9) in the conditions of Theorem 1 using (2.2) and passing to the limit (1.11) in (2.3).

Let us take some λ_0, $\rho(\lambda_0)>0$, where ρ is defined in (1.4). Putting in formula (2.3) $\lambda=\lambda_0+\xi/n$ and $\mu=\lambda_0+\eta/n$, we get:

$$\begin{aligned}&\frac{1}{n}K_n(\lambda_0+\xi/n,\lambda_0+\eta/n)=\\&\qquad-\frac{c_{m,n}}{4\pi^2}\oint\limits_{L}\oint\limits_{\omega}\frac{\exp\{m((\lambda_0+\xi/n)u-(\lambda_0+\eta/n)t)\}}{u-t}\frac{t^m}{u^m}\prod_{j=1}^{n}\frac{u-\tau_j}{t-\tau_j}d\,u\,d\,t\\&\qquad=-\frac{c_{m,n}}{4\pi^2}\oint\limits_{L}\oint\limits_{\omega}\frac{\exp\{m(S_n(u,\lambda_0)-S_n(t,\lambda_0))\}}{(u-t)}e^{(\xi u-\eta t)c_{m,n}}d\,u\,d\,t,\end{aligned}\tag{3.1}$$ $\boxed{\texttt{Ker1}}$

where

$$S_n(z,\lambda_0)=\lambda_0 z-\ln z+\frac{c_{m,n}^{-1}}{n}\sum_{j=1}^{n}\ln(z-\tau_j)-S^*,\quad c_{m,n}=\frac{m}{n},\quad \tau_j=1/t_j^{(n)}, \tag{3.2}$$ $\boxed{\texttt{S\_n}}$

and S^* is a constant which will be chosen later (see ($\overset{\texttt{S*}}{3.20}$)). Here L is a closed contour, encircling $\{\tau_j\}_{j=1}^n$ and ω is any closed contour encircling 0 and not intersecting L. Thus, ($\overset{\texttt{Ker1}}{3.1}$) and ($\overset{\texttt{S\_n}}{3.2}$) yield

$$\frac{1}{n}K_n(\lambda_0+\xi/n,\lambda_0+\eta/n)=-\frac{c_{m,n}}{4\pi^2}\oint\limits_L\oint\limits_\omega \mathcal{F}_n(t,u)d\,u\,d\,t, \tag{3.3}$$ $\boxed{\texttt{Ker}}$

where to simplify formulas below we denote

$$\mathcal{F}_n(t,u)=\frac{\exp\{m(S_n(u,\lambda_0)-S_n(t,\lambda_0))\}}{(u-t)}e^{(\xi u-\eta t)c_{m,n}}. \tag{3.4}$$ $\boxed{\texttt{F\_cal}}$

We start from the following statement

$\boxed{\texttt{p:rav\_sh}}$ **Proposition 3.** *Set*

$$g_n^{(0)}=\frac{1}{n}\sum\frac{1}{\tau_j-z},\quad f^{(0)}=\int\frac{N^{(0)}(d\,t)}{\tau-z},\quad \tau=1/t. \tag{3.5}$$ $\boxed{\texttt{g\_0,f\_0}}$

Then we have under conditions of Theorem $\overset{\texttt{thm:1}}{1}$

$$\lim_{n\to\infty}\mathbf{P}\{|g_n^{(0)}(z)-f^{(0)}(z)|>\varepsilon\}=0 \tag{3.6}$$ $\boxed{\texttt{rav\_sh}}$

uniformly in z from compact set K in the upper half-plane. Moreover, the inverse assertion is true, i.e., if ($\overset{\texttt{rav\_sh}}{3.6}$) is valid for some compact set K in the upper half-plane, then we have ($\overset{\texttt{conpN0}}{1.14}$).

The proof of the proposition is given, e.g., in $\overset{\texttt{TSh:08}}{[10]}$.

Let us take the disk $\sigma=\{z:\,|z_0-z|\le\varepsilon_1\}$ as the compact set K, where

$$z_0=-c^{-1}f(\lambda_0-i0)-\frac{1-c^{-1}}{\lambda_0} \tag{3.7}$$ $\boxed{\texttt{z\_0}}$

with f defined in ($\overset{\texttt{eqv\_f}}{1.13}$). Here we note that according to the result of $\overset{\texttt{Sil-Ch:95}}{[9]}$ $f(\lambda_0-i0)$ exists and it is a continuous function for $\lambda_0>0$ and

$$\Im z_0=-c^{-1}\Im f(\lambda_0-i0)=c^{-1}\pi\rho(\lambda_0)>0. \tag{3.8}$$ $\boxed{\texttt{z\_0\_rho}}$

Taking into account ($\overset{\texttt{rav\_sh}}{3.6}$), we get that for any sufficiently small $\delta>0$ and for any $\varepsilon>0$ there exists n_0 such that for all $n>n_0$ the event

$$\Omega_\varepsilon=\{\max_{z\in\sigma}|g_n^{(0)}(z)-f^{(0)}(z)|<\varepsilon\}, \tag{3.9}$$ $\boxed{\texttt{Om\_e}}$

satisfies the condition

$$\mathbf{P}_n^{(\tau)}\{\Omega_\varepsilon^C\}\le\delta.$$

Recall that $\mathbf{P}_n^{(\tau)}$ and $\mathbf{E}_n^{(\tau)}$ correspond to the averaging with respect to Σ_n.

According to the determinant formulas (2.2)-(2.3), we have to prove that for any $k \in \mathbb{N}$

$$\lim_{n\to\infty} \mathbf{E}_n^{(\tau)} \left\{ \det \left\{ \frac{1}{n\rho_n(\lambda_0)} K_n\left(\lambda_0 + \xi_i/n\rho_n(\lambda_0), \lambda_0 + \xi_j/n\rho_n(\lambda_0)\right) \right\}_{i,j=1}^{k} \right\} = \det\left\{S(\xi_i - \xi_j)\right\}_{i,j=1}^{k}, \tag{3.10}$$

where S is define in (1.6). Consider the expression

$$\mathbf{E}_n^{(\tau)} \left\{ \det \left\{ \frac{1}{n\rho_n(\lambda_0)} K_n\left(\lambda_0 + \xi_i/n\rho_n(\lambda_0), \lambda_0 + \xi_j/n\rho_n(\lambda_0)\right) \right\}_{i,j=1}^{k} - \det\left\{S(\xi_i - \xi_j)\right\}_{i,j=1}^{k} \right\}. \tag{3.11}$$

It is easy to see that to obtain (1.5) it suffices to prove that

$$\begin{aligned} \left| \frac{1}{n\rho_n(\lambda_0)} K_n\left(\lambda_0 + \xi/n\rho_n(\lambda_0), \lambda_0 + \eta/n\rho_n(\lambda_0)\right) - \alpha(\xi,\eta) S(\xi - \eta) \right| &\le C\varepsilon, \quad \text{on } \Omega_\varepsilon, \\ \left| \frac{1}{n\rho_n(\lambda_0)} K_n\left(\lambda_0 + \xi/n\rho_n(\lambda_0), \lambda_0 + \eta/n\rho_n(\lambda_0)\right) \right| &\le C, \qquad \text{on } \Omega_\varepsilon^C. \end{aligned} \tag{3.12}$$

where S and K_n are defined in (1.6) and (3.3) respectively and $\alpha(\xi,\eta)$ is some multiplier which vanishes during the calculation of the determinant in (2.2). Indeed, choose sufficiently small ε and δ and split $\mathbf{E}_n^{(\tau)}\{\ldots\}$ in (3.11) into two parts, the integral over Ω_ε of (3.9) and the integral over its complement. If we know (3.12), then the first integral is small because the difference of the determinants is small on Ω_ε. In view of the Hadamard inequality ([20], Section I.5) and the second line of (3.12), we have

$$\begin{aligned} &\left| \det \left\{ \frac{1}{n\rho_n(\lambda_0)} K_n\left(\lambda_0 + \xi_i/n\rho_n(\lambda_0), \lambda_0 + \xi_j/n\rho_n(\lambda_0)\right) \right\}_{i,j=1}^{k} \right| \\ &\qquad \le \prod_{i=1}^{k} \left(\sum_{j=1}^{k} \left| \frac{1}{n\rho_n(\lambda_0)} K_n\left(\lambda_0 + \xi_i/n\rho_n(\lambda_0), \lambda_0 + \xi_j/n\rho_n(\lambda_0)\right) \right|^2 \right)^{1/2} \le k^{k/2} C^k. \end{aligned} \tag{3.13}$$

Besides, $\det\{S(\xi_i - \xi_j)\}_{i,j=1}^{k}$, with S of (1.6), is bounded. Thus, the integral over Ω_ε^C in (3.11) is bounded by $C\delta$ and hence we obtain that (3.11) is bounded by $C\varepsilon_1$. Thus, we are left to prove (3.12).

To do this we will choose the contour L in (3.3) as some n-dependent contour L_n. Consider the equation for z with a real parameter $\lambda > 0$

$$V(z) := \frac{1}{z} + c_{m,n}^{-1} g_n^{(0)}(z) = \lambda, \tag{3.14}$$

where $g_n^{(0)}$ is defined in (3.5). Equation (3.14) can be written as a polynomial equation of degree $(n+1)$ (if $\lambda > 0$) and so it has $(n+1)$ roots. Consider $z \in \mathbb{R}$. If $z \to \tau_j + 0$, then $V(z) \to -\infty$, if $z \to \tau_j - 0$, then $V(z) \to +\infty$, and also if $z \to +0$, then $V(z) \to +\infty$, if $z \to -0$, then $V(z) \to -\infty$. Besides, $V(z) \to +0$, as $z \to +\infty$ (since $c_{m,n} > 1$) and $V(z) \to -0$, as $z \to -\infty$. Thus, the graph of $V(z)$ for $z \in \mathbb{R}$ looks like in Fig.1.



$n-1$ roots of (3.14) are always real and lying between τ_j-s. If λ is big enough, then all $n+1$ roots are distinct and real. Let $z_n(\lambda)$ be a root which tends to 0, as $\lambda \to \infty$. If λ decreases, then at some $\lambda = \lambda_{c_1}$ two roots coincide and for $\lambda \le \lambda_{c_1}$ the real root disappears and there appear two complex roots – $z_n(\lambda)$ and $\overline{z_n(\lambda)}$. Indeed, it is easy to see, that since (3.14) has not more than $n+1$ roots, these coinciding roots are positive and smaller than any τ_j. Thus $\lambda_{c_1} > 0$, since for $z^* = z_n(\lambda_{c_1})$ we have

$$V(z^*) = \frac{1}{z_*} + \frac{c_{m,n}^{-1}}{n} \sum_{j=1}^{n} \frac{1}{\tau_j - z_*} > 0,$$

because $z_* > 0$ and $\tau_j > z_*$ for $j = 1,..,n$. Then $z_n(\lambda)$ may be real again, than again complex, and so on, however, as soon as λ becomes less then some $\lambda_{c_2} > 0$, $z_n(\lambda)$ becomes again real (and it is real for any $0 < \lambda < \lambda_{c_2}$). Moreover, $z_n(\lambda_{c_2})$ is bigger that every τ_j. Choose

$$\begin{aligned} L_n = \{z \in \mathbb{C} : z = z_n(\lambda) :\ \Im z_n(\lambda) > 0,\ \lambda > 0\} \\ \cup \{z \in \mathbb{C} : z = \overline{z_n(\lambda)} :\ \Im z_n(\lambda) > 0,\ \lambda > 0\} \cup S', \end{aligned} \tag{3.15}$$

where S' is a set of points $z = z_n(\lambda)$ in which $z_n(\lambda)$ becomes real. Since (3.14) always has exactly $n+1$ roots, it is clear that the set of λ's corresponding to $z_n(\lambda) \in L_n$ is $\bigcup_{j=1}^{k} I_k$, where $\{I_j\}_{j=1}^{k}$ are non intersecting segments. It is easy to see also that the contour L_n is closed and encircling $\{\tau_j\}_{j=1}^{n}$, and L_n lies in the right half plane. In addition, we will use later that for real z $V'(z)$ can change the sign only in the points of S'.

To prove (3.12), let us prove first that

$$\begin{aligned} \left| \frac{1}{n} K_n\left(\lambda_0 + \xi/n, \lambda_0 + \eta/n\right) - \alpha(\xi,\eta) \frac{\sin\left(\pi c_{m,n} y_n(\lambda_0)(\eta - \xi)\right)}{\pi(\xi - \eta)} \right| \le C\varepsilon, \quad \text{on } \Omega_\varepsilon, \\ \left| \frac{1}{n} K_n\left(\lambda_0 + \xi/n, \lambda_0 + \eta/n\right) \right| \le C, \qquad\qquad \text{on } \Omega_\varepsilon^C, \end{aligned} \tag{3.16}$$

where $y_n(\lambda) = \Im z_n(\lambda)$, K_n and Ω_ε are defined in (3.3) and (3.9), $z_n(\lambda)$ is a root of (3.14) which tends to $+0$, as $\lambda \to \infty$, and $\alpha(\xi,\eta)$ is some multiplier which vanishes during the computation of the determinant in (2.2).

To do this we replace the integration over ω by the integration over ω_n, where

$$\omega_n = \{z \in \mathbb{C} : z = u(\varphi) = re^{i\varphi},\ \varphi \in [0; 2\pi)\} \tag{3.17}$$ om_n

with $r = |z_n(\lambda_0)|$. Consider the contour ω_δ of Fig.2, where δ is small enough. It will be shown below (see Lemmas 1, 2) that L_n and ω_n has only two points of intersection: $z_n(\lambda_0)$ and $\overline{z_n(\lambda_0)}$.



According to the residue theorem, we have

$$\oint\limits_{\omega_\delta} \mathcal{F}_n(t,u)d\,u = \begin{cases} 2\pi i \cdot \exp\{(\xi-\eta)tc_{m,n}\}, & t \text{ is inside } \omega_\delta, \\ 0, & t \text{ is outside } \omega_\delta, \\ \pi i \cdot \exp\{(\xi-\eta)tc_{m,n}\}, & t = z_n(\lambda_0), \overline{z_n(\lambda_0)}, \end{cases}$$

where $\mathcal{F}_n$ is defined in (3.4). The sum of the integrals over the lines $\Im z = \pm\delta$ tends to 0, as $\delta \to 0$, hence we get after the limit $\delta \to 0$

$$\begin{aligned} \oint\limits_{L_n} \oint\limits_{\omega \cup \omega_n} \mathcal{F}_n(t,u)d\,u\,d\,t &= 2\pi i \int\limits_{z_n(\lambda_0)}^{\overline{z_n(\lambda_0)}} \exp\{(\xi-\eta)tc_{m,n}\}\,d\,t \\ &= 4\pi c_{m,n}^{-1} e^{x_n(\lambda_0)(\xi-\eta)c_{m,n}} \frac{\sin((\xi-\eta)y_n(\lambda_0)c_{m,n})}{\xi-\eta}. \end{aligned} \tag{3.18}$$ Res

Taking into account that

$$\oint\limits_{L_n}\oint\limits_{\omega} \mathcal{F}_n(t,u)d\,u\,d\,t = \oint\limits_{L_n}\oint\limits_{\omega_n} \mathcal{F}_n(t,u)d\,u\,d\,t - \oint\limits_{L_n}\oint\limits_{\omega \cup \omega_n} \mathcal{F}_n(t,u)d\,u\,d\,t$$

and since the r.h.s. of (3.18) is bounded (see Lemma 3 below), to prove (3.16) it suffices to prove that

$$\oint\limits_{L_n}\oint\limits_{\omega_n} |\mathcal{F}_n(t,u)|\,d\,u\,d\,t \le \begin{cases} C\varepsilon, & \text{on } \Omega_\varepsilon, \\ C, & \text{on } \Omega_\varepsilon^C, \end{cases} \tag{3.19}$$ lim0

where $\mathcal{F}_n(t,u)$ is defined in (3.4).

Now let us choose the constant in (3.2) as

$$S^* = \Re\left(\lambda_0 z_n(\lambda_0) - \ln z_n(\lambda_0) + \frac{c_{m,n}^{-1}}{n}\sum_{j=1}^{n}\ln(z_n(\lambda_0)-\tau_j)\right) \tag{3.20}$$ `S*`

and study the behavior of the function $\Re S_n(z_n(\lambda),\lambda_0)$ of (3.2) on the contours L_n of (3.15) and ω_n of (3.17).

`l:min_L` **Lemma 1.** *Let $z \in L_n$ of (3.15) and $\Im z \ge 0$. Then for any set $\{\tau_j\}_{j=1}^n \in (\mathbb{R}^+)^n$ we have*

$$\Re S_n(z_n(\lambda),\lambda_0) \ge 0$$

and the equality holds only at $\lambda = \lambda_0$. Moreover, the function $\Re S_n(z_n(\lambda),\lambda_0)$ is strictly increasing for $\lambda > \lambda_0$ and strictly decreasing for $\lambda < \lambda_0$. The same is valid for the lower part of L_n, i.e., $z \in L_n$, $\Im z < 0$.

`l:max_om` **Lemma 2.** *Consider $u \in \omega_n$, $\Im u > 0$. Then we have for any set $\{\tau_j\}_{j=1}^n \in (\mathbb{R}^+)^n$*

$$\Re S_n(u,\lambda_0) \le -C|x-x_0|^2,$$

where $x_0 = \Re z_n(\lambda_0)$, $x = \Re u$. The same is valid for the lower part of ω_n.

These lemmas yield that for $t \in L_n$, $u \in \omega_n$ and for any set $\{\tau_j\}_{j=1}^n \in (\mathbb{R}^+)^n$

$$\Re((S_n(u,\lambda_0) - S_n(t,\lambda_0))) \le 0,$$

and the equality holds only if u and t are both equal to $z_n(\lambda_0)$ or $\overline{z_n(\lambda_0)}$.

To estimate $\mathcal{F}_n(t,u)$ of (3.4) we use also

`l:prexp` **Lemma 3.** *There exists an n-independent $\delta > 0$ such that for any $\lambda \in U_\delta(\lambda_0)$ uniformly in $\{\tau_j\}_{j=1}^n \in (\mathbb{R}^+)^n$ the solution $z_n(\lambda)$ of equation (3.14) admits the following bounds*

$$0 < C_1 < x_n(\lambda) < C_2, \quad 0 < C_1 < |z_n(\lambda)| < C_2. \tag{3.21}$$ `preexp`

where $x_n(\lambda) = \Re z_n(\lambda)$ (here and below $U_\delta(a) = (a-\delta, a+\delta)$).

In particular, this means that for any set $\{\tau_j\}_{j=1}^n \in (\mathbb{R}^+)^n$ the radius r of ω_n of (3.17) satisfies the inequality

$$0 < C_1 < r < C_2. \tag{3.22}$$ `ineqv_r`

Now we split the integral in (3.19) into two parts

$$\oint_{L_n}\oint_{\omega_n} |\mathcal{F}_n(t,u)|\,d\,u\,d\,t = \left(\oint_{\omega_n}\int_{L_n^A} + \oint_{\omega_n}\int_{L_n\setminus L_n^A}\right)|\mathcal{F}_n(t,u)|d\,u\,d\,t \tag{3.23}$$ `Int_sk`

where L_n^A is the part of L_n where $|x_n(\lambda)| \le A$, $x_n(\lambda) = \Re z_n(\lambda)$ and $\mathcal{F}_n(t,u)$ is defined in (3.4).

The next lemma implies the bound for the length of L_n

`l:dl_kont` **Lemma 4.** *Let $l(x)$ be the oriented length of L_n between $x_0 = x_n(\lambda_0)$ and x ($l(x) \ge 0$ for $x > x_0$). Then uniformly in $\{\tau_j\}_{j=1}^n \in (\mathbb{R}^+)^n$ $l(x)$ admits the bound:*

$$|l(x_1) - l(x_2)| \le C_1|x_1 - x_2| + C_2(|x_1| + |x_2|) + C_3.$$

Moreover, if $0 < x_1 < x_2 < C$, $j = 1, 2$, then

$$l(x_2) - l(x_1) \le C\sqrt{x_2 - x_1}.$$

To estimate the r.h.s. of (3.23) we use

`l:int_okr` **Lemma 5.** *There exists a constant $\delta > 0$ such that for all $\{\tau_j\}_{j=1}^n \subset (\mathbb{R}^+)^n$ and any $x \notin U_\delta(x_0)$ we have*

$$dist\{z_n(x), \omega_n\} \geq \delta,$$

where $z_n(x)$ is $z_n(\lambda)$ which is expressed via $x_n(\lambda)$ (we can do it according to (3.35)).

Moreover, if $L_{[x_1,x_2]}$ is a part of L_n between lines $\Re z = x_1$ and $\Re z = x_2$, $x_0 = \Re z_n(\lambda_0)$ and $\mu^* = \min\limits_{z \in L_{[x_1,x_2]}} \Re S_n(z, \lambda_0)$ (recall that $\mu^* \geq 0$). Then for any $\{\tau_j\}_{j=1}^n \in (\mathbb{R}^+)^n$

$$\int\limits_{L_{[x_1,x_2]}} \oint\limits_{\omega_n} |\mathcal{F}_n(t,u)|\, d\,u\, d\,t \leq \begin{cases} Ce^{-m\mu^*}, & 0 < x_1 < x_2 \leq x_0/2, \\ Ce^{-m\mu^*}\sqrt{x_2 - x_1}\,|\ln(x_2 - x_1)|, & x_0/2 < x_1 < x_2 < C. \end{cases} \tag{3.24}$$ `L_otr`

Thus, the first integral in (3.23) is bounded by $C = C(A)$ (we split L_n^A in two segments by the point $x = x_0/2$ and take the sum of the bounds for this segments).

To prove that the second integral in (3.23) is bounded we consider the imaginary part of (3.14). We get

$$\frac{c_{m,n}^{-1}}{n} \sum_{j=1}^n \frac{1}{(x_n(\lambda) - \tau_j)^2 + y_n^2(\lambda)} = \frac{1}{x_n^2(\lambda) + y_n^2(\lambda)}, \tag{3.25}$$ `cond`

where $x_n(\lambda) = \Re z_n(\lambda)$, $y_n(\lambda) = \Im z_n(\lambda)$. This and (3.21) yield for $|z_n(\lambda)| > A$

$$\begin{aligned} \frac{1}{n}\sum_{j=1}^n (\ln|z_n(\lambda_0) - \tau_j| - \ln|z_n(\lambda) - \tau_j|) &= \frac{1}{n}\sum_{j=1}^n \ln\left|1 + \frac{z_n(\lambda_0) - z_n(\lambda)}{z_n(\lambda) - \tau_j}\right| \\ \leq \frac{1}{n}\sum_{j=1}^n \frac{|z_n(\lambda_0) - z_n(\lambda)|}{|z_n(\lambda) - \tau_j|} &\leq |z_n(\lambda_0) - z_n(\lambda)| \left(\frac{1}{n}\sum_{j=1}^n \frac{1}{|z_n(\lambda) - \tau_j|^2}\right)^{1/2} \\ &\leq \frac{c_{m,n}^{1/2}|z_n(\lambda_0) - z_n(\lambda)|}{|z_n(\lambda)|} \leq C. \end{aligned} \tag{3.26}$$ `razn_log`

Using (3.20), (3.26) and Lemma 3, we obtain for $z_n \in L_n \setminus L_n^A$

$$\begin{aligned} \Re S_n(z_n(\lambda), \lambda_0) &= \lambda_0 x_n(\lambda) - \ln|z_n(\lambda)| + \frac{c_{m,n}^{-1}}{n}\sum_{j=1}^n \ln|z_n(\lambda) - \tau_j| - S^* \\ &\geq \lambda_0 x_n(\lambda) - \frac{1}{2}\ln|z_n(\lambda)|^2 + C. \end{aligned} \tag{3.27}$$ `ots_S_vsp`

Moreover, it follows from (3.25) that there exists j such that

$$y_n^{-2}(\lambda) \geq \frac{1}{(x_n(\lambda) - \tau_j)^2 + y_n^2(\lambda)} \geq \frac{c_{m,n}}{x_n^2(\lambda) + y_n^2(\lambda)},$$

and hence

$$(c_{m,n} - 1)y_n^2(\lambda) \leq x_n^2(\lambda). \tag{3.28}$$ `ineqv_y`

Therefore, (3.27) yields

$$\Re S_n(z_n(\lambda), \lambda_0) \geq \lambda_0 x_n(\lambda) - C_1 \ln|x_n(\lambda)| + C_2 \geq \frac{\lambda_0 x_n(\lambda)}{2} + C, \tag{3.29}$$ `ots_S`

if $|x_n(\lambda)| > A$, where $A \in \mathbb{N}$ is big enough, but independent of $\{\tau_j\}_{j=1}^n$ and n. Besides, Lemma 3 yields for sufficiently big A

$$\frac{1}{|u-t|} \le \frac{1}{|\Re u - \Re t|} \le \frac{1}{A-r} \le C.$$

This, Lemmas 2-3, (3.22) and (3.29) imply (recall that $m/n = c_{m,n}$)

$$\int\limits_{L_n \setminus L_n^A} \oint\limits_{\omega_n} |\mathcal{F}_n(t,u)|\, d\,u\, d\,t \le C_1 \int\limits_A^\infty e^{-nxC_2 + C_3 n} d\,l(x),$$

where $l(x)$ is defined in Lemma 4 and $C_2 > 0$. According to Lemma 4 we obtain for sufficiently big A and n

$$\begin{aligned}\int\limits_A^\infty e^{-nC_2 x + C_3 n} d\,l(x) &= \sum_{k=A}^\infty \int\limits_k^{k+1} e^{-nC_2 x + nC_3} d\,l(x)\\ \le \sum_{k=A}^\infty e^{-nkC_2 + nC_3}(l(k+1) - l(k)) &\le \sum_{k=A}^\infty e^{-nkC_2 + nC_3}(C_3 + C_4 k) < e^{-nd}\end{aligned}$$

for some $d > 0$. Thus,

$$\int\limits_{L_n \setminus L_n^A} \oint\limits_{\omega_n} |\mathcal{F}_n(t,u)|\, d\,u\, d\,t \le e^{-nd}, \qquad (3.30) \quad \texttt{int\_A}$$

and we have proved the second line of (3.19) (see (3.23), Lemma 5, and (3.30)).

Let us prove now the first line of (3.19). Choose any sufficiently small $\delta > 0$ and split the integral in (3.19) into three parts

$$\oint\limits_{L_n} \oint\limits_{\omega_n} |\mathcal{F}_n(t,u)|\, d\,u\, d\,t = \left(\oint\limits_{\omega_n} \int\limits_{U_1} + \oint\limits_{\omega_n} \int\limits_{L_n^A \setminus U_1} + \oint\limits_{\omega_n} \int\limits_{L_n \setminus L_n^A} \right) |\mathcal{F}_n(t,u)| d\,u\, d\,t \qquad (3.31) \quad \texttt{Int\_sk1}$$

where $U_1 = \{z_n(\lambda), \overline{z_n(\lambda)} : \lambda \in U_\delta(\lambda_0)\}$ and L_n^A is defined in (3.23). Use

`l:x'` **Lemma 6.** *For $\{\tau_j\}_{j=1}^n \in \Omega_\varepsilon$ and for $\lambda \in U_\delta(\lambda_0)$ with sufficiently small δ we have*

$$0 < C_1 < |x_n'(\lambda)| < C_2.$$

This lemma and Lemma 5 yield

$$\oint\limits_{\omega_n} \int\limits_{U_1} |\mathcal{F}_n(t,u)|\, d\,u\, d\,t \le \oint\limits_{\omega_n} \int\limits_{L_{\delta'}} |\mathcal{F}_n(t,u)| \le C\sqrt{\delta} \ln \delta^{-1},$$

where U_1 is defined in (3.31), $\delta' = O(\delta)$, $L_{\delta'}$ is a part of L_n between the lines $\Re z = x_n(\lambda_0) + \delta'$ and $\Re z = x_n(\lambda_0) - \delta'$ and δ is small enough.

To estimates the second integral in (3.31) we need more information about the behavior of $\Re S_n(z, \lambda_0)$ on L_n. Consider the second derivative of $\Re S_n(z, \lambda_0)$. We have

`l:vt_pr` **Lemma 7.** *There exist $C > 0$ and $\delta_0 > 0$ such that for $\{\tau_j\}_{j=1}^n \in \Omega_\varepsilon$*

$$\frac{d^2}{d\,\lambda^2} \Re(-S_n(z_n(\lambda), \lambda_0)) < -C$$

for any $\lambda \in U_{\delta_0}(\lambda_0)$.

According to Lemma 7 and the equalities

$$\Re S_n(z_n(\lambda_0),\lambda_0)=0,\quad \frac{d}{d\,\lambda}S_n(z_n(\lambda),\lambda_0)\Big|_{\lambda=\lambda_0}=0,$$

(see (3.2), (3.20), and (3.34)) we obtain for any $\delta<\delta_0$

$$\Re(-S_n(z_n(\lambda),\lambda_0))<-C\frac{(\lambda-\lambda_0)^2}{2},\ \lambda\in U_\delta(\lambda_0).$$

Since the function $\Re\left(S_n(z_n(\lambda),\lambda_0)\right)$ is monotone for $\lambda\neq\lambda_0$ (see Lemma 1), we get

$$\Re(-S_n(z_n(\lambda),\lambda_0))<-C\frac{\delta^2}{2},\quad \lambda\notin U_\delta(\lambda_0).$$

This and Lemmas 4-6 yield

$$\oint\limits_{\omega_n}\int\limits_{L_n^A\setminus U_1}|\mathcal{F}_n(t,u)|\,d\,u\,d\,t\leq C\exp\{-nC\delta^2/2\}.$$

The bound for the last integral in (3.31) is obtained in (3.30). Therefore, we have proved (3.19), and combining it with (3.3) and (3.18), we get

$$\left|\frac{1}{n}K_n\left(\lambda_0+\xi/n,\lambda_0+\eta/n\right)-e^{(\xi-\eta)x_n(\lambda_0)c_{m,n}}\frac{\sin\left((\xi-\eta)y_n(\lambda_0)c_{m,n}\right)}{\pi(\xi-\eta)}\right|\leq\begin{cases}C\varepsilon, & \text{on }\Omega_\varepsilon,\\ C, & \text{on }\Omega_\varepsilon^C.\end{cases}\tag{3.32}$$ Main

Thus we proved the first line of (3.16). The second line of (3.16) follows from (3.32), Lemma 3 and the bound

$$\frac{\sin\left((\xi-\eta)y_n(\lambda_0)c_{m,n}\right)}{\pi(\xi-\eta)}\leq\pi^{-1}c_{m,n}y_n(\lambda_0).$$

Therefore, (3.16) is proved.

To prove (3.12) we are left to prove that $y_n(\lambda_0)c_{m,n}=\rho_n(\lambda_0)+o(1)$, $n\to\infty$. Consider the limiting equation

$$\frac{1}{z}+c^{-1}f^{(0)}(z)=\lambda,\tag{3.33}$$ eqv_f_0

where $f^{(0)}$ is defined in (3.5) and $\lambda\in\mathbb{R}$ is fixed.

l:8 **Lemma 8.** *There exists ε' such that for any $\lambda\in U_{\varepsilon'}(\lambda_0)$ equation (3.33) has a unique solution $z(\lambda)$ in the upper half-plane $\Im z>0$, moreover $z(\lambda_0)=z_0$, where z_0 is defined in (3.7). The solution $z(\lambda)$ is continuous in $\lambda\in U_{\varepsilon'}(\lambda_0)$ and $\Im z(\lambda_0)>0$.*

Relation between $z(\lambda)$ and $z_n(\lambda)$ is given by

l:9 **Lemma 9.** *There exists δ such that for all $\lambda\in U_\delta(\lambda_0)$ and for sufficiently big n*

$$|z_n(\lambda)-z(\lambda)|<\varepsilon\quad \textit{on }\Omega_\varepsilon.$$

The determinant formulas (2.2)-(2.3) imply that $\rho_n(\lambda)=\frac{1}{n}\mathbf{E}_n^{(\tau)}\{K_n(\lambda,\lambda)\}$, where ρ_n, K_n is defined in (1.2) and (2.3). Then

$$\begin{aligned}&\rho_n(\lambda_0)-\rho(\lambda_0)\\&\qquad=\mathbf{E}_n^{(\tau)}\left\{(n^{-1}K_n(\lambda_0,\lambda_0)-\rho(\lambda_0))\mathbf{1}_{\Omega_\varepsilon}\right\}+\mathbf{E}_n^{(\tau)}\left\{(n^{-1}K_n(\lambda_0,\lambda_0)-\rho(\lambda_0))\mathbf{1}_{\Omega_\varepsilon^C}\right\}.\end{aligned}$$

According to the second line of (3.16), the second term can be bounded by $C\delta$. According to Lemma 9, $|z_n(\lambda_0) - z(\lambda_0)| < \varepsilon_1$, where $z_n(\lambda_0)$ and $z(\lambda_0)$ are the solutions of equation (3.14) and (3.33) for $\lambda = \lambda_0$. Therefore, using Lemma 3 and (3.8), we have

$$\left|\pi^{-1} c y_n(\lambda_0) - \rho(\lambda_0)\right| < \varepsilon_1,$$

where $y_n(\lambda) = \Im z_n(\lambda)$. This and the first line of (3.32) for $\xi = \eta = 0$ yield on Ω_ε for any $\varepsilon_1 > 0$

$$|n^{-1}K_n(\lambda_0,\lambda_0) - \rho(\lambda_0)| < |n^{-1}K_n(\lambda_0,\lambda_0) - \pi^{-1}cy_n(\lambda_0)| + |\pi^{-1}cy_n(\lambda_0) - \rho(\lambda_0)| \le 2\varepsilon_1$$

as $m, n \to \infty$. Therefore, for any $\varepsilon_1 > 0$ there exists such N that we have for any $n > N$

$$|\rho_n(\lambda_0) - \rho(\lambda_0)| < C\varepsilon_1.$$

Thus, $\rho_n(\lambda_0) > 0$ for sufficiently big n and we can divide by it in (3.16). Therefore, we obtain (3.12) and thus (1.5).

Let us prove (1.8). It is well-known (see,e.g.,[11]) that

$$\begin{aligned} &E_n\left([\lambda_0 + a/\rho_n(\lambda_0)n, \lambda_0 + b/\rho_n(\lambda_0)n]\right) \\ &\qquad = 1 + \sum_{l=1}^{\infty} \frac{(-1)^l}{l!} \int_a^b \det\left\{n^{-1}K_n\left(\lambda_0 + x_i/\rho_n(\lambda_0)\, n, \lambda_0 + x_j/\rho_n(\lambda_0)\, n\right)\right\}_{i,j=1}^{l} \prod_{j=1}^{l} d\, x_j. \end{aligned}$$

Thus, according to the dominant convergence theorem, (1.5) and (3.13) yield (1.8). Therefore, Theorem 1 is proved.

3.1 Proofs of the lemmas

Proof of Lemma 1. Differentiate $\Re S_n(z_n(\lambda), \lambda_0)$ with respect to λ. Using (3.2) and equation (3.14), we obtain

$$\begin{aligned} \frac{d}{d\,\lambda}\Re S_n(z_n(\lambda),\lambda_0) &= \Re\left(z_n'(\lambda)\left(\lambda_0 - \frac{1}{z_n(\lambda)} - c_{m,n}^{-1} g_n^{(0)}(z_n(\lambda))\right)\right) \\ &= \Re\, z_n'(\lambda)\,(\lambda_0 - \lambda) = -x_n'(\lambda)(\lambda - \lambda_0). \end{aligned} \tag{3.34}$$

Let us show now that

$$x_n'(\lambda) < 0. \tag{3.35}$$

Differentiating (3.14) with respect to λ we get

$$z_n'(\lambda) = \left(c_{m,n}^{-1} \frac{d}{d\,z} g_n^{(0)}(z_n(\lambda)) - \frac{1}{z_n^2(\lambda)}\right)^{-1}. \tag{3.36}$$

It follows from the implicit function theorem that L_n intersects the real axis at the points where

$$c_{m,n}^{-1} \frac{d}{d\,x} g_n^{(0)}(x) - \frac{1}{x^2} = 0.$$

Since

$$\frac{d}{d\,x} g_n^{(0)}(x) = \frac{1}{n}\sum_{j=1}^{n} \frac{1}{(x-\tau_j)^2},$$

the inequality $c_{m,n}^{-1}\dfrac{d}{d\,x}g_n^{(0)}(x)-\dfrac{1}{x^2}>0$ holds near τ_j. As we note before, $V'(x)$ change sign only at the points of S' in which L_n intersects real axis. Thus, the function $c_{m,n}^{-1}\dfrac{d}{d\,x}g_n^{(0)}(x)-\dfrac{1}{x^2}$ is always negative outside L_n. On the other hand, $z_n(\lambda)=x_n(\lambda)$ for $x>0$ outside L_n and in this case

$$x_n'(\lambda)=z_n'(\lambda)=\left(c_{m,n}^{-1}\frac{d}{d\,z}g_n^{(0)}(z_n(\lambda))-\frac{1}{z_n^2(\lambda)}\right)^{-1}<0.$$

Let now $\lambda\in\bigcup\limits_{j=1}^{k}I_j$, i.e., $z_n(\lambda)$ belongs to L_n. We get from ($\overset{\texttt{zn\_pr}}{3.36}$)

$$\Re z_n'(\lambda)=x_n'(\lambda)=\frac{a_n(\lambda)}{a_n^2(\lambda)+b_n^2(\lambda)},\quad \Im z_n'(\lambda)=y_n'(\lambda)=-\frac{b_n(\lambda)}{a_n^2(\lambda)+b_n^2(\lambda)},$$

where

$$\begin{aligned} a_n(\lambda)\ &=\Re\left(c_{m,n}^{-1}\frac{d}{d\,z}g_n^{(0)}(z_n(\lambda))-\frac{1}{z_n^2(\lambda)}\right),\\ b_n(\lambda)\ &=\Im\left(c_{m,n}^{-1}\frac{d}{d\,z}g_n^{(0)}(z_n(\lambda))-\frac{1}{z_n^2(\lambda)}\right).\end{aligned}\qquad(3.37)\quad\boxed{\texttt{ab}}$$

Using ($\overset{\texttt{cond}}{3.25}$), we obtain

$$\begin{aligned} a_n(\lambda)\ &=\ \frac{c_{m,n}^{-1}}{n}\sum_{j=1}^{n}\frac{(x_n(\lambda)-\tau_j)^2-y_n^2(\lambda)}{|z_n(\lambda)-\tau_j|^4}-\frac{x_n^2(\lambda)-y_n^2(\lambda)}{|z_n(\lambda)|^4}\\ &=\ \frac{2y_n^2(\lambda)}{|z_n(\lambda)|^4}-\frac{c_{m,n}^{-1}}{n}\sum_{j=1}^{n}\frac{2y_n^2(\lambda)}{|z_n(\lambda)-\tau_j|^4}.\end{aligned}\qquad(3.38)$$

Besides, according ($\overset{\texttt{cond}}{3.25}$) we have

$$\frac{1}{n}\sum_{j=1}^{n}\frac{1}{|z_n(\lambda)-\tau_j|^4}\geq\left(\frac{1}{n}\sum_{j=1}^{n}\frac{1}{|z_n(\lambda)-\tau_j|^2}\right)^2=\frac{c_{m,n}^2}{|z_\lambda|^4}.$$

Then, since $c_{m,n}\to c>1$, as $m,n\to\infty$, ($\overset{\texttt{a}}{3}$.38) yields

$$a_n(\lambda)<-2y_n^2(\lambda)\frac{c_{m,n}-1}{|z_n(\lambda)|^4}<0\qquad(3.39)\quad\boxed{\texttt{ineqv2}}$$

Thus, in this case we also have

$$x_n'(\lambda)<0.\qquad(3.40)\quad\boxed{\texttt{Rezpr}}$$

According to ($\overset{\texttt{S1}}{3}$.34), this means that the function $\Re S_n(z_n(\lambda),\lambda_0)$ is strictly increasing for $\lambda>\lambda_0$ and strictly decreasing for $\lambda<\lambda_0$, i.e. $\Re S_n(z,\lambda_0)$ has a minimum at $\lambda=\lambda_0$. Since $\Re S_n(z_n(\lambda_0),\lambda_0)=0$, $\Re S_n(z_n(\lambda),\lambda_0)\geq 0$ and the equality holds only at $\lambda=\lambda_0$.

Note that the lower part of L_n differs from the upper one only by the sign of $y_n(\lambda)$, hence $\Re S_n(z,\lambda_0)\geq 0$, $z\in L_n$ and the equality holds only at $z=z_n(\lambda_0)$ and $z=\overline{z_n(\lambda_0)}$. $\square$

Proof of the Lemma $\overset{\texttt{l:max\_om}}{\mathbf{2}}$. $\Re S_n(z,\lambda_0)$ on ω_n of ($\overset{\texttt{om\_n}}{3.17}$) we can rewrite as

$$\Re S_n(u(\varphi),\lambda_0)=\lambda_0 r\cos\varphi-\ln r+\frac{c_{m,n}^{-1}}{2n}\sum_{j=1}^{n}\ln(r^2-2r\tau_j\cos\varphi+\tau_j^2)-S^*.$$

Differentiating this with respect to $\varphi \in (0, \pi)$ we obtain

$$\Re S_n(u(\varphi), \lambda_0)' = r \sin\varphi \left(\frac{c_{m,n}^{-1}}{n} \sum_{j=1}^{n} \frac{\tau_j}{r^2 - 2r\tau_j \cos\varphi + \tau_j^2} - \lambda_0 \right). \tag{3.41}$$ `Su1`

Set

$$\varphi_0 = \arg\, z_n(\lambda_0).$$

It is easy to see (using (`cond`3.25) and (`eqv_g_0_n`3.14)), that the expression in the brackets in (`Su1`3.41) is zero at $\varphi = \varphi_0$, and, moreover, this expression is a strictly monotone decreasing function of φ. Thus $\varphi = \varphi_0$ is a maximum of $\Re S_n(u(\varphi), \lambda_0)$. Similarly the point $\varphi = -\varphi_0$ is a maximum for lower-half plane.

Besides, using that $r = |z_n(\lambda_0)|$, we have for $u \in \omega_n$

$$\Re S_n(u, \lambda_0) = \lambda_0 \widetilde{x} - \ln r + \frac{c_{m,n}^{-1}}{2n} \sum_{j=1}^{n} \ln(r^2 - 2\widetilde{x}\tau_j + \tau_j^2) - S^*,$$

where $\widetilde{x} = \Re u$. Thus,

$$\frac{d}{d\,\widetilde{x}} \Re S_n(u, \lambda_0) = \lambda_0 - \frac{c_{m,n}^{-1}}{n} \sum_{j=1}^{n} \frac{\tau_j}{r^2 - 2\widetilde{x}\tau_j + \tau_j^2}.$$

Differentiating with respect to $\widetilde{x}$ twice we get for $u \in \omega_n$, $\widetilde{x} = \Re u$

$$\begin{aligned} \frac{d^2}{d\,\widetilde{x}^2} \Re S_n(u, \lambda_0) &= -\frac{2c_{m,n}^{-1}}{n} \sum_{j=1}^{n} \frac{\tau_j^2}{(r^2 - 2\widetilde{x}\tau_j + \tau_j^2)^2} \le -2c_{m,n}^{-1} \left(\frac{1}{n} \sum_{j=1}^{n} \frac{\tau_j}{r^2 - 2\widetilde{x}\tau_j + \tau_j^2} \right)^2 \\ &= -2c_{m,n} \left(\frac{d}{d\,\widetilde{x}} \Re S_n(u, \lambda_0) - \lambda_0 \right)^2 \le 4c_{m,n}\lambda_0 \frac{d}{d\,\widetilde{x}} \Re S_n(u, \lambda_0) - 2c_{m,n}\lambda_0^2. \end{aligned}$$

Since we proved that $\widetilde{x} = x_0 = \Re z_n(\lambda_0)$ is a maximum $\Re S_n(u, \lambda_0)$ on ω_n, where $x_0 = \Re z_n(\lambda_0)$, and $\Re S_n(z_n(\lambda_0), \lambda_0) = 0$, integrating from x_0 to $\widetilde{x} > x_0$ we obtain

$$\frac{d}{d\,\widetilde{x}} \Re S_n(u, \lambda_0) \le 4c_{m,n}\lambda_0 \Re S_n(u, \lambda_0) - 2c_{m,n}\lambda_0^2(\widetilde{x} - x_0),$$

where $\widetilde{x} = \Re u$. Hence, the Gronuol lemma yields for $\widetilde{x} - x_0 > 0$

$$\Re S_n(u, \lambda_0) \le \frac{\lambda_0}{2}(\widetilde{x} - x_0) + \frac{c_{m,n}^{-1}}{8} \left(1 - e^{4c_{m,n}\lambda_0(\widetilde{x} - x_0)} \right) \le -C(\widetilde{x} - x_0)^2,$$

where $u \in \omega_n$, $\widetilde{x} = \Re u$. It is easy to see, that similar inequality we have for $\widetilde{x} - x_0 < 0$ and therefore we get

$$\Re S_n(u, \lambda_0) \le -C(\Re u - x_0)^2$$

for any $u \in \omega_n$. Since the lower part of ω_n differs from the upper one only by the sign of $\Im u$, the proof is complete. □

Proof of Lemma `l:prexp`3. Equation (`eqv_g_0_n`3.14) has $(n+1)$ roots $z_1, \ldots, z_{n+1}$ for $\lambda \ne 0$. According to the Viet theorem

$$z_1 + z_2 + \ldots + z_{n+1} = \frac{1 - c_{m,n}^{-1}}{\lambda} + \sum_{j=1}^{n} \tau_j.$$

Moreover, $n-1$ of these roots are always real and lying to the left of the corresponding τ_j, and hence their sum is less than $\sum_{j=1}^{n} \tau_j$ (see Fig.1). Therefore, since we consider $\lambda > 0$, the sum of the last two roots are greater than $(1-c_{m,n}^{-1})/\lambda$.

If the last two roots of (3.14) are real (i.e. $y_n(\lambda) = 0$), then we obtain

$$x_n'(\lambda) = \left(\frac{c_{m,n}^{-1}}{n} \sum_{j=1}^{n} \frac{1}{(x_n(\lambda)-\tau_j)^2} - \frac{1}{x_n^2(\lambda)} \right)^{-1}.$$

According to (3.35), we get

$$0 < \frac{1}{x_n^2(\lambda)} - \frac{c_{m,n}^{-1}}{n} \sum_{j=1}^{n} \frac{1}{(x_n(\lambda)-\tau_j)^2} < \frac{1}{x_n^2(\lambda)} - c_{m,n}^{-1} \left(\frac{1}{n} \sum_{j=1}^{n} \frac{1}{x_n(\lambda)-\tau_j} \right)^2$$
$$= \frac{1}{x_n^2(\lambda)} - c_{m,n} \left(\frac{1}{x_n(\lambda)} - \lambda \right)^2.$$

Therefore,

$$\frac{1-c_{m,n}^{-1/2}}{\lambda} \le x_n(\lambda) \le \frac{1+c_{m,n}^{-1/2}}{\lambda}$$

and for $\lambda > \delta > 0$ such that $y_n(\lambda) = 0$ the lemma is proved.

If the last two roots of (3.14) are complex ($z_n(\lambda)$ and $\overline{z_n(\lambda)}$) and $y_n(\lambda) \neq 0$, then

$$\frac{1-c_{m,n}^{-1}}{\lambda} < z_n(\lambda) + \overline{z_n(\lambda)} = 2x_n(\lambda).$$

Moreover, we know from (3.14) that

$$\frac{1}{z_n(\lambda)} = \lambda + \frac{c_{m,n}^{-1}}{n} \sum_{j=1}^{n} \frac{1}{z_n(\lambda)-\tau_j}.$$

The real part of the l.h.s. of this equation is less than $\frac{1}{x_n(\lambda)}$. On the other hand, the real part of the r.h.s. is greater than $\lambda - \frac{c_{m,n}^{-1}}{n} \sum_{j=1}^{n} \frac{|x_n(\lambda)-\tau_j|}{|z_n(\lambda)-\tau_j|^2}$. But for $z_n(\lambda)$ such that $y_n(\lambda) \neq 0$, (3.25) yields

$$\frac{1}{n} \sum_{j=1}^{n} \frac{|x_n(\lambda)-\tau_j|}{|z_n(\lambda)-\tau_j|^2} \le \left(\frac{1}{n} \sum_{j=1}^{n} \frac{|x_n(\lambda)-\tau_j|^2}{|z_n(\lambda)-\tau_j|^2} \right)^{1/2} \left(\frac{1}{n} \sum_{j=1}^{n} \frac{1}{|z_n(\lambda)-\tau_j|^2} \right)^{1/2}$$
$$\le \frac{\sqrt{c_{m,n}}}{|z_n(\lambda)|} \le \frac{\sqrt{c_{m,n}}}{|x_n(\lambda)|}.$$

Hence, for $\lambda > \delta > 0$

$$\frac{1+c_{m,n}^{-1/2}}{|x_n(\lambda)|} \ge \lambda \Rightarrow |x_n(\lambda)| \le \frac{1+c_{m,n}^{-1/2}}{\lambda} < C,$$

which complete the proof of the first inequality of (3.21) for $y_n(\lambda) \neq 0$ (i.e. $z_n(\lambda) \in L_n$).

The second inequality in (3.21) for $\lambda : y_n(\lambda) \neq 0$ easily follows from the first inequality and (3.28). The lemma is proved. $\square$

Proof of Lemma `l:dl_kont` **4.** It follows from (`Rezpr` 3.40) and (`a` 3.38) that one can express $y_n(\lambda)$ via $x_n(\lambda)$ to obtain the "graph" $y_n(x)$ of the upper part of L_n. Denote

$$y_n^2(x) = s(x), \quad x - \tau_j = \triangle_j, \quad \sigma_0 = \frac{1}{x^2+s}$$
$$\sigma_k = \frac{1}{n}\sum_{j=1}^{n}\frac{1}{(\triangle_j^2+s)^k}, \quad \sigma_{kl} = \frac{1}{n}\sum_{j=1}^{n}\frac{\triangle_j^l}{(\triangle_j^2+s)^k}, \ (k=\overline{1,3},\ l=1,2) \tag{3.42}$$ `obozn`

Differentiating (`cond` 3.25) with respect to x, we obtain the equality

$$s'(x)\left(c_{m,n}^{-1}\sigma_2 - \sigma_0^2\right) + 2c_{m,n}^{-1}\sigma_{21} - 2x\sigma_0^2 = 0.$$

Note that

$$x\sigma_0^2 \le \sigma_0^{3/2}, \quad \sigma_2 \ge \sigma_1^2 = c_{m,n}^2\sigma_0^2$$

$$\sigma_{21} \le \left(\frac{1}{n}\sum_{j=1}^{n}\frac{\triangle_j^2}{(\triangle_j^2+s(x))^2}\right)^{1/2}\left(\frac{1}{n}\sum_{j=1}^{n}\frac{1}{(\triangle_j^2+s(x))^2}\right)^{1/2} = \sigma_{22}^{1/2}\sigma_2^{1/2} \le (c_{m,n}\sigma_0\sigma_2)^{1/2},$$

thus

$$\begin{aligned}(s'(x))^2 = \frac{4(x\sigma_0^2 - c_{m,n}^{-1}\sigma_{21})^2}{(c_{m,n}^{-1}\sigma_2-\sigma_0^2)^2} &\le 8\left(\frac{x^2\sigma_0^4}{(c_{m,n}^{-1}\sigma_2-\sigma_0^2)^2} + \frac{c_{m,n}^{-2}\sigma_{21}^2}{(c_{m,n}^{-1}\sigma_2-\sigma_0^2)^2}\right)\\ &\le 8\left(\frac{1}{(c_{m,n}-1)^2\sigma_0} + \frac{c_{m,n}^{-1}\sigma_2}{(c_{m,n}-1)\sigma_0(c_{m,n}^{-1}\sigma_2-\sigma_0^2)}\right).\end{aligned} \tag{3.43}$$ `s1_1a`

Since

$$\frac{c_{m,n}^{-1}\sigma_2}{c_{m,n}^{-1}\sigma_2-\sigma_0^2} = 1 + \frac{\sigma_0^2}{c_{m,n}^{-1}\sigma_2-\sigma_0^2} \le 1 + \frac{1}{c_{m,n}-1},$$

(`s1_1a` 3.43) yields for $(x, y(x)) \in L_n$

$$(s'(x))^2 \le \frac{C}{\sigma_0} = C(x^2+y^2(x)).$$

Thus, according to Lemma `l:prexp` 3 we obtain for any x such that $(x, y(x)) \in L_n$ and $|x| < C$

$$|s'(x)| \le C. \tag{3.44}$$ `s'`

Differentiating (`cond` 3.25) with respect to x twice, we have in our notations

$$\begin{aligned}s''(x)(c_{m,n}^{-1}\sigma_2-\sigma_0^2) &= 2(s'(x))^2(c_{m,n}^{-1}\sigma_3-\sigma_0^3)\\ &\quad + 8s'(x)(c_{m,n}^{-1}\sigma_{31}-x\sigma_0^3) + 6(c_{m,n}^{-1}\sigma_2-\sigma_0^2) + 8s(x)(\sigma_0^3 - c_{m,n}^{-1}\sigma_3).\end{aligned}$$

Since $\sigma_3 \ge (c_{m,n}\sigma_0)^3 > c_{m,n}\sigma_0^3$, the r.h.s. is a quadratic trinomial of $s'(x)$ with a positive leading coefficient (without addition with $s(x)$). Hence, using $c_{m,n}^{-1}\sigma_2 - \sigma_0^2 > 0$ we get

$$\begin{aligned}s''(x)(c_{m,n}^{-1}\sigma_2-\sigma_0^2) &\ge 8s(x)(\sigma_0^3 - c_{m,n}^{-1}\sigma_3) - \frac{8(c_{m,n}^{-1}\sigma_{31}-x\sigma_0^3)^2}{c_{m,n}^{-1}\sigma_3-\sigma_0^3}\\ &\ge -8s(x)c_{m,n}^{-1}\sigma_3 - \frac{16c_{m,n}^{-2}\sigma_{31}^2}{c_{m,n}^{-1}\sigma_3-\sigma_0^3} - \frac{16x^2\sigma_0^6}{c_{m,n}^{-1}\sigma_3-\sigma_0^3}.\end{aligned} \tag{3.45}$$ `s2_1`

Note that $s(x)\sigma_3 \le \sigma_2$ and also

$$\sigma_{31}^2 = \left(\frac{1}{n}\sum_{j=1}^{n} \frac{\triangle_j}{(\triangle_j^2 + s(x))^3}\right)^2 \le \sigma_{23}\sigma_3 \le \sigma_2\sigma_3.$$

Using these inequalities, we get from (`s2_1`3.45)

$$s''(x)(c_{m,n}^{-1}\sigma_2 - \sigma_0^2) \ge -8c_{m,n}^{-1}\sigma_2 - \frac{16c_{m,n}^{-2}\sigma_2\sigma_3}{c_{m,n}^{-1}\sigma_3 - \sigma_0^3} - \frac{16\sigma_0^5}{c_{m,n}^{-1}\sigma_3 - \sigma_0^3}.$$

Thus, since

$$c_{m,n}^{-1}\sigma_2 - \sigma_0^2 \ge (c_{m,n} - 1)\sigma_0^2 > 0, \quad c_{m,n}^{-1}\sigma_3 - \sigma_0^3 \ge (c_{m,n}^2 - 1)\sigma_0^3,$$

$$\frac{c_{m,n}^{-1}\sigma_2}{c_{m,n}^{-1}\sigma_2 - \sigma_0^2} = 1 + \frac{\sigma_0^2}{c_{m,n}^{-1}\sigma_2 - \sigma_0^2} \le \frac{c_{m,n}}{c_{m,n} - 1},$$
$$\frac{c_{m,n}^{-1}\sigma_3}{c_{m,n}^{-1}\sigma_3 - \sigma_0^3} = 1 + \frac{\sigma_0^3}{c_{m,n}^{-1}\sigma_3 - \sigma_0^3} \le \frac{c_{m,n}^2}{c_{m,n}^2 - 1},$$

we obtain

$$s''(x) \ge -C. \tag{3.46}$$
`s2`

Let $y_n'(x) = \frac{s'(x)}{2\sqrt{s(x)}} > 0$ when $x \in [x_1, x_2]$. Then we have

$$l(x_2) - l(x_1) = \int\limits_{x_1}^{x_2} \sqrt{1 + (y_n'(x))^2}\,d\,x = \int\limits_{x_1}^{x_2} \sqrt{1 + \left(\frac{s'(x)}{2\sqrt{s(x)}}\right)^2}\,d\,x$$
$$\le \int\limits_{x_1}^{x_2} \left(1 + \frac{s'(x)}{2\sqrt{s(x)}}\right) d\,x = (x_2 - x_1) + \sqrt{s_2} - \sqrt{s_1} \le (x_2 - x_1) + \sqrt{s_2 - s_1}, \tag{3.47}$$
`ots_l1`

where $s_2 = s(x_2)$, $s_1 = s(x_1)$. If we choose in (`ots_l1`3.47) x_2 being the maximum point of $s(x)$, then $s'(x_2) = 0$ and we can write

$$s_1 - s_2 = \frac{s''(\zeta)(x_1 - x_2)^2}{2},$$

where $\zeta \in [x_1, x_2]$. This and (`s2`3.46) imply

$$0 \le s_2 - s_1 \le C(x_1 - x_2)^2.$$

Hence, we get in view of (`ots_l1`3.47)

$$l(x_2) - l(x_1) \le C(x_2 - x_1). \tag{3.48}$$
`ots_l2`

We have similar inequality for $x_1 > x_2$ and $y_n'(x) < 0$, $x \in [x_2, x_1]$ (we should consider $l(x_1) - l(x_2)$). Take an arbitrary $x_1 \in [x_0; x_2]$ and denote by x_* and x^* the nearest to x_1 and x_2 extremal points of $s(x)$ in $[x_1, x_2]$ correspondingly. Then, splitting $[x_*, x_*]$ in the segments of monotonicity of y_n and using (`ots_l1`3.47), its analog for decreasing $y_n(x)$ and (`ots_l2`3.48) we obtain

$$\begin{aligned} l(x_2) - l(x_1) &= l(x_*) - l(x_1) + l(x^*) - l(x_*) + l(x_2) - l(x^*) \\ &\le C(x^* - x_*) + (x_2 - x^*) + \sqrt{|s_2 - s^*|} + (x_* - x_1) + \sqrt{|s_* - s_1|}, \end{aligned} \tag{3.49}$$
`ots_l_kor`

where Let now x_1, x_2 be such that $|x_1|, |x_2| \le C$. Then (3.49) and (3.44) yield

$$l(x_2) - l(x_1) \le C(x_2 - x_1) + C_1\sqrt{x_2 - x^*} + \sqrt{x_* - x_1} \le C\sqrt{x_2 - x_1},$$

and so the second statement of the lemma is proved.

Also it easy to see from (3.47) or its analog for decreasing $y_n(x)$ that for any segment $[a, b]$ where $y_n(x)$ is monotone we have

$$l(b) - l(a) \le |b - a| + |y_n(b) - y_n(a)|. \qquad (3.50) \quad \boxed{\texttt{ots\_l3}}$$

Again take an arbitrary $x_1 \in [x_0; x_2]$ and denote by x_* and x^* the nearest to x_1 and x_2 extremal points of $s(x)$ in $[x_1, x_2]$ correspondingly. Using (3.48), its analog for decreasing $y_n(x)$, and (3.50) we obtain

$$\begin{aligned} l(x_2) - l(x_1) &= l(x_*) - l(x_1) + l(x^*) - l(x_*) + l(x_2) - l(x^*) \\ &\le C(x_2 - x_0) + |y_n(x_2) - y_n(x^*)| + |y_n(x_1) - y_n(x_*)|. \end{aligned} \qquad (3.51)$$

Taking into account (3.28) it is easy to see that

$$y_n(x) \le C_1|x| + C_2.$$

This and (3.51) yield

$$|l(x_2) - l(x_1)| \le C_1|x_2 - x_1| + C_2(|x_1| + |x_2|) + C_3.$$

□

Proof of Lemma 5. Note that since r is bounded from both sides uniformly in $\{\tau_j\}_{j=1}^n$ and n (see (3.22)), we have for $|z| \le C_1/2$ (here C_1 is a constant from (3.22))

$$|z - u(\varphi)| \ge r - C_1/2,$$

and the assertions of the lemma follow from Lemmas 1-2 and 4. If $z \in L_n$, $|z| > C_1/2$ then similarly to (3.26) we get

$$\begin{aligned} \Re(S_n(z,\lambda_0) - S_n(u(\varphi),\lambda_0)) &= \lambda_0(z - u(\varphi)) + \ln|u(\varphi)| - \ln|z| \\ + \frac{c_{m,n}^{-1}}{n}\sum_{j=1}^n \ln|z_n(\lambda) - \tau_j| &- \frac{c_{m,n}^{-1}}{n}\sum_{j=1}^n \ln|u(\varphi) - \tau_j| \le \frac{C_2|z_n(\lambda) - u(\varphi)|}{|\Im u(\varphi)|} \end{aligned}$$

This inequality, Lemmas 1, 2 yield for $z \in L_n$, $|z| > C_1/2$ and $u(\varphi) \in \omega_n$

$$C|\Re u(\varphi) - x_0|^2 \le -\Re S_n(u(\varphi),\lambda_0) \le \Re(S_n(z,\lambda_0) - S_n(u(\varphi),\lambda_0)) \le \frac{C_2|z - u(\varphi)|}{|\Im u(\varphi)|}.$$

Hence, we obtain

$$|z - u(\varphi)| \ge Cr|\Re u(\varphi) - x_0|^2 \sin\varphi, \quad z \in L_n, \quad |z| > C_1/2, \quad \varphi \in [0, \pi/2]. \qquad (3.52) \quad \boxed{\texttt{in\_cu\_1}}$$

Thus for any $u(\varphi) \in \omega_n$ there are no points of L_n in the disk of radius $Cr|\Re u(\varphi) - x_0|^2 \sin\varphi$ centered in $u(\varphi)$. Consider two curves (see Fig.3):

$$\begin{aligned} &\widetilde{z}_\pm(\varphi) = r(1 \pm C|\Re u(\varphi) - x_0|^2 \sin\varphi)e^{i\varphi}, \quad \varphi \in [0, \pi/2) \\ &\text{or} \quad \begin{cases} \widetilde{x}_\pm(\varphi) = r(\cos\varphi \pm C|\Re u(\varphi) - x_0|^2 \sin\varphi\cos\varphi), \\ \widetilde{y}_\pm(\varphi) = r(\sin\varphi \pm C|\Re u(\varphi) - x_0|^2 \sin^2\varphi), \end{cases} \end{aligned} \qquad (3.53) \quad \boxed{\texttt{curv}}$$

where $x_0 = \Re z_n(\lambda_0)$.



It is easy to see $\widetilde{x}'_+(\varphi) < 0$, $\widetilde{y}'_+(\varphi) \geq 0$ when φ is closed to π, and $\widetilde{x}'_+(\varphi) > 0$, $\widetilde{y}'_+(\varphi) \geq 0$ when φ is closed to $\varphi_0 = \arg z_n(\lambda_0)$. Let φ^* be the nearest to 0 root of $\widetilde{x}'_+(\varphi)$ on $[0, \pi/2]$, and $x^* = \widetilde{x}'_+(\varphi^*)$. It is evident that $x^* - r > C^*$, where C^* is a constant depending only on the curve $\widetilde{z}_+(\varphi)$ and therefore bounded from below uniformly in $\{\tau_j\}_{j=1}^n$ and n. Moreover, there are no points of L_n in the domain D^* bounded by the curve $\widetilde{z}_+(\varphi)$ and lines $\Im z = 0$ and $\Re z = x^*$. Indeed, since we know, that L_n does not intersect $\widetilde{z}_+(\varphi)$, to come from $z_n(\lambda_0)$ to any point of D^*, $z_n(\lambda)$ should intersect the line $\Re z = x^*$ twice (or more), that contradict to (3.35). Now taking into account that for $\lambda \to \infty$ $z_n(\lambda)$ is inside of ω_n and therefore for $x > x_0$ $z_n(x)$ is outside of ω_n, we obtain the first assertion of the lemma.

Now, if $[x_1, x_2] \cap U_\delta(x_0) = \emptyset$, then (3.24) follows from Lemmas 1-2 and from Lemma 4. If $[x_1, x_2] \subset U_\delta(x_0)$, let us write

$$\int\limits_{L_{[x_1,x_2]}} \oint\limits_{\omega_n} |\mathcal{F}_n(u,t)| d\,u\,d\,t \leq Ce^{-m\mu*} \int\limits_{L_{[x_1,x_2]}} \oint\limits_{\omega_n} \frac{|d\,u|\,|d\,t|}{|u-t|}$$

Using that $|t| \geq x_0/2$ for $t \in L_{[x_1,x_2]}$, $x_1 > x_0/2$, we get

$$\oint\limits_{\omega_n} \frac{|d\,u|}{|u-t|} = r\int\limits_0^{2\pi} \frac{d\,\varphi}{\sqrt{r^2 - 2r|t|\cos\varphi + |t|^2}} = \int\limits_0^{2\pi} \frac{d\,\varphi}{\sqrt{(r-|t|)^2 + 4r|t|\sin^2\dfrac{\varphi}{2}}}$$

$$\leq 2\sqrt{2}\int\limits_0^{\pi} \frac{d\,\varphi}{|r-|t|| + 2\sqrt{r|t|}\sin\frac{\varphi}{2}} \leq C(1 + \ln|r-|t||^{-1}),$$

where $\varphi = \arg(u-t)$ Hence to prove the second statement of the Lemma 5 we should bound the integral of $\ln|r-|t||^{-1}$. To this end we use the curves $\widetilde{y}_\pm(x)$. Note that $\widetilde{y}'_\pm(\varphi) > 0$ and $\widetilde{x}'_\pm(\varphi) > 0$ for $\varphi \in U_{\delta_1}(\varphi_0)$. Hence, we can choose sufficiently small δ (independent on $\{\tau_j\}_{j=1}^n$ and n) such that both curves $\widetilde{z}_\pm(\varphi)$ can be parameterized by $x \in U_\delta(x_0)$. Then according to the above argument we have

$$\begin{cases} y_n(x) \leq \widetilde{y}_-(x), & x \leq x_0, \quad x \in U_\delta(x_0), \\ y_n(x) \geq \widetilde{y}_+(x), & x \geq x_0, \quad x \in U_\delta(x_0). \end{cases} \tag{3.54}$$

Note that since $\cos\varphi, \sin\varphi > C_0 > 0$ for $\widetilde{x}_{\pm}(\varphi) \in U_\delta(x_0)$, we get

$$\begin{aligned}|\widetilde{x}_{\pm}(\varphi) - r\cos\varphi| &\geq C|r\cos\varphi - x_0|^2 = C|r\cos\varphi - \widetilde{x}_{\pm}(\varphi) + \widetilde{x}_{\pm}(\varphi) - x_0|^2 \\ &\geq C(|\widetilde{x}_{\pm}(\varphi) - x_0|^2 - 2|\widetilde{x}_{\pm}(\varphi) - x_0| \cdot |r\cos\varphi - \widetilde{x}_{\pm}(\varphi)|).\end{aligned}$$

Therefore,

$$|\widetilde{z}_{\pm}(\varphi) - u(\varphi)| \geq |\widetilde{x}_{\pm}(\varphi) - r\cos\varphi| \geq C|\widetilde{x}_{\pm}(\varphi) - x_0|^2. \tag{3.55} \quad \boxed{\texttt{in1}}$$

Thus, for $x_0 - \delta \leq x_1 < x_2 \leq x_0$, using (3.54)-(3.55) we obtain

$$\begin{aligned}I(x_1, x_2) &:= -\int_{x_1}^{x_2} \ln\left(r - \sqrt{x^2 + y_n^2(x)}\right) d\,l(x) \leq -\int_{x_1}^{x_2} \ln\left(r - \sqrt{x^2 + \widetilde{y}_-^2(x)}\right) d\,l(x) \\ &= -\int_{x_1}^{x_2} \ln|u(\varphi_-(x)) - \widetilde{z}_-(x)|\, d\,l(x) \leq -\int_{x_1}^{x_2} \ln\left(C|x - x_0|^2\right) d\,l(x).\end{aligned} \tag{3.56} \quad \boxed{\texttt{in2}}$$

Besides, Lemma 4 implies that for any $x_1 < x_2 \leq x_0$ we have

$$\begin{aligned}&-\int_{x_1}^{x_2} \ln|x - x_0| d\,l(x) = -(l(x_1) - l(x_2)) \ln|x_1 - x_0| - \int_{x_1}^{x_2} \frac{l(x) - l(x_2)}{x_0 - x} d\,x \\ &\leq C\sqrt{x_2 - x_1}|\ln(x_2 - x_1)| + C\int_{x_1}^{x_2} \frac{\sqrt{x_2 - x}}{x_2 - x} d\,x \leq C\sqrt{x_2 - x_1}|\ln(x_2 - x_1)|.\end{aligned} \tag{3.57} \quad \boxed{\texttt{b\_int\_g}}$$

For $x_0 \leq x_1 < x_2 \leq x_0 + \delta$ the proof is similar, we need only use that $y_n(x) \geq \widetilde{y}_+(x)$ in (3.56). Now it is straightforward to obtain the statement of Lemma 5 for any $x_1, x_2 > x_0/2$ (if it is necessary, we split $[x_1, x_2]$ on the segments by the points $x_0 - \delta$ and $x_0 + \delta$ and then take the sum of the bounds for these segments). $\square$

Proof of Lemma 8. The existence of $f(\lambda_0 \pm i0)$ is proved in [9]. Using (1.13) we obtain

$$\frac{1}{z(\lambda_0)} + c^{-1} f^{(0)}(z(\lambda_0)) = \lambda_0, \tag{3.58} \quad \boxed{\texttt{eqv\_l0}}$$

where $z(\lambda_0)$ is defined in (3.7). Hence, the solution exists if $\lambda = \lambda_0$. Besides, for $x(\lambda_0) = \Re z(\lambda_0)$, $y(\lambda_0) = \Re z(\lambda_0)$ we have

$$\begin{aligned}&\Re\left(\frac{1}{z} + c^{-1} f^{(0)}(z)\right)'_z \Bigg|_{z=z(\lambda_0)} \\ &\qquad = c^{-1}\int \frac{((x(\lambda_0) - \tau)^2 - y^2(\lambda_0))\, N^{(0)}(d\,t)}{((x(\lambda_0) - \tau)^2 + y^2(\lambda_0))^2} - \frac{(x^2(\lambda_0) - y^2(\lambda_0))}{(x^2(\lambda_0) + y^2(\lambda_0))^2},\end{aligned} \tag{3.59} \quad \boxed{\texttt{pr\_z}}$$

where $\tau = 1/t$. Since $y(\lambda_0) \neq 0$, considering the l.h.s. of (3.58) we get

$$\frac{1}{x^2(\lambda_0) + y^2(\lambda_0)} = c^{-1}\int \frac{N^{(0)}(d\,t)}{(x(\lambda_0) - \tau)^2 + y^2(\lambda_0)}.$$

Thus, we can rewrite (3.59) as

$$\begin{aligned}&\Re\left(\frac{1}{z} + c^{-1} f^{(0)}(z)\right)'_z \Bigg|_{z=z(\lambda_0)} \\ &\qquad = -\left(\int \frac{2c^{-1} y^2(\lambda_0) N^{(0)}(d\,t)}{((x(\lambda_0) - \tau)^2 + y^2(\lambda_0))^2} - \frac{2y^2(\lambda_0)}{(x^2(\lambda_0) + y^2(\lambda_0))^2}\right),\end{aligned} \tag{3.60} \quad \boxed{\texttt{pr\_z1}}$$

where $\tau = 1/t$. But we know that

$$\int \frac{N^{(0)}(d\,t)}{((x(\lambda_0)-\tau)^2+y^2(\lambda_0))^2} \geq \left(\int \frac{N^{(0)}(d\,t)}{(x(\lambda_0)-\tau)^2+y^2(\lambda_0)}\right)^2 = \frac{c^2}{(x^2(\lambda_0)+y^2(\lambda_0))^2}.$$

Therefore, since $c > 1$, (3.60) yields

$$\Re\left(\frac{1}{z}+c^{-1}f^{(0)}(z)\right)'_z\Bigg|_{z=z(\lambda_0)} < -\frac{2y^2(\lambda_0)(c-1)}{(x^2(\lambda_0)+y^2(\lambda_0))^2} < 0.$$

Hence, according to the implicit function theorem, equation (3.33) has a unique solution in the upper half-plane $\Im z > 0$ if $\lambda \in U_\varepsilon(\lambda_0)$, moreover the solution is continuous in λ in this neighborhood. $\square$

Proof of the Lemma 7. We get from (3.34)

$$\frac{d^2}{d\,\lambda^2}\Re(-S_n(z_n(\lambda),\lambda_0)) = \frac{d}{d\,\lambda}\left(x'_n(\lambda)(\lambda-\lambda_0)\right) = x'_n(\lambda)+x''_n(\lambda)(\lambda-\lambda_0). \qquad (3.61)$$

Hence, to prove the lemma it is sufficient to show that $x''_n(\lambda)$ is bounded uniformly in n and that $x'_n(\lambda)$ is bounded from below by a positive constant uniformly in n in some sufficiently small neighborhood $U_\delta(\lambda_0)$.

Show that $-x'_n(\lambda) \geq C_1$ for all $\lambda \in U_\delta(\lambda_0)$, where C_1 is a positive n-independent constant. We have from (3.36)

$$-\Re z'_n(\lambda) = -x'_n(\lambda) = \frac{-a_n(\lambda)}{a_n^2(\lambda)+b_n^2(\lambda)},$$

where a_n, b_n are defined in (3.37). Let us prove that $a_n(\lambda)$ and $b_n(\lambda)$ are bounded uniformly in n and that $a_n(\lambda)$ is bounded from below by a positive constant uniformly in n for all $\lambda \in U_\delta(\lambda_0)$ (evidently, it will be sufficient).

Note that for $z_n(\lambda) \in L_n$

$$|b_n(\lambda)| = \left|\frac{2y_n(\lambda)x_n(\lambda)}{|z_n(\lambda)|^4} - \frac{c_{m,n}^{-1}}{n}\sum_{j=1}^{n}\frac{2y_n(\lambda)(x_n(\lambda)-\tau_j)}{|z_n(\lambda)-\tau_j|^4}\right| \leq \frac{2|y_n(\lambda)||x_n(\lambda)|}{|z_n(\lambda)|^4}$$
$$+\frac{c_{m,n}^{-1}}{n}\sum_{j=1}^{n}\frac{2|y_n(\lambda)||x_n(\lambda)-\tau_j|}{|z_n(\lambda)-\tau_j|^4} \leq \frac{1}{|z_n(\lambda)|^2}+\frac{c_{m,n}^{-1}}{n}\sum_{j=1}^{n}\frac{1}{|z_n(\lambda)-\tau_j|^2} = \frac{2}{|z_\lambda|^2}$$

Also we can see from (3.38) that

$$|a_n(\lambda)| \leq \left|\frac{2y_n^2(\lambda)}{|z_n(\lambda)|^4}+\frac{c_{m,n}^{-1}}{n}\sum_{j=1}^{n}\frac{2y_n^2(\lambda)}{|z_n(\lambda)-\tau_j|^4}\right| \leq \frac{4}{|z_n(\lambda)|^2},$$

and besides (3.39) yields

$$-a_n(\lambda) > 2y_n^2(\lambda)\frac{c_{m,n}-1}{|z_n(\lambda)|^4}.$$

Using Lemma 9, continuity of $z(\lambda)$, and since $\Im z(\lambda_0) = \pi\rho(\lambda_0) > 0$, we have starting from some n

$$|x_n(\lambda)| < C_1, \quad |y_n(\lambda)| < C_1, \quad |y_n(\lambda)| > C_2 \qquad (3.62)$$

for all $\lambda \in U_\delta(\lambda_0)$, where n-independent δ is small enough.

This fact yields that $-x'_n(\lambda) \geq C$ for all $\lambda \in U_\delta(\lambda_0)$, where C is a positive n-independent constant, and also that x''_n is bounded uniformly and hence the second terms in (3.61) is of order δ. The lemma is proved. $\square$

Proof of Lemma 9. We use Lemma 8. Consider the solution $z(\lambda)$ of the limiting equation (3.33) in some neighborhood of λ_0. Since $\lambda_0 \in \operatorname{supp} N$, $\Im z(\lambda_0) = A > 0$. Taking into account the continuity of $z(\lambda)$ near λ_0, we can take a sufficiently small neighborhood $U_{\delta_1}(\lambda_0)$ such that

$$|z(\lambda) - z(\lambda_0)| < \varepsilon/2, \ \lambda \in U_{\delta_1}(\lambda_0).$$

Consider the set of the functions $f_\lambda(z) = \dfrac{1}{z} + c^{-1} f^{(0)}(z) - \lambda$ and the function

$$\phi(z) = c_{m,n}^{-1} g_n^{(0)}(z) - c^{-1} f^{(0)}(z),$$

where $f^{(0)}$, $g_n^{(0)}$ are defined in (3.5), and set $\omega = \{z : |z - z(\lambda_0)| \le \varepsilon\}$. Let us show that for any $\lambda \in U_{\delta_1}(\lambda_0)$ and for any $z \in \omega$

$$|f_\lambda(z)| > c_0, \tag{3.63}$$

where c_0 does not depend on λ. Assume the opposite and choose a sequence $\{\lambda_k\}_{k \ge 1}$, $\lambda_k \in U_{\delta_1}(\lambda_0)$ such that $|f_{\lambda_k}(z_k)| \to 0$, as $k \to \infty$. There exists a subsequence $\{\lambda_{k_l}\}$, converging to some $\lambda \in U_{\delta_1}(\lambda_0)$ such that the subsequence $\{z_{k_l}\}$ converges to $z \in \partial\omega$. For these λ and z $f_\lambda(z) = 0$. But equation $f_\lambda(z) = 0$ has in the upper half-plane only one root $z(\lambda)$, which is inside of the circle of the radius $\varepsilon/2$ and with the center $z(\lambda_0)$. This contradiction proves (3.63). Since $|g_n^{(0)}(z) - f^{(0)}(z)| \le \varepsilon$ on Ω_ε uniformly on ω (see (3.9) and $c_{m,n} \to c$, we have starting from some n

$$|\phi(z)| < c_0, \ z \in \partial\omega. \tag{3.64}$$

Comparing (3.63) and (3.64), we obtain that starting from some n

$$|f_\lambda(z)| > |\phi(z)|, \ z \in \partial\omega, \ \forall \lambda \in U_{\delta_1}(\lambda_0).$$

Since both functions are analytic, the Rouchet theorem implies that $f_\lambda(z)$ and $f_\lambda(z) + \phi(z) = \dfrac{1}{z} + c_{m,n}^{-1} g_n^{(0)}(z) - \lambda$ have the same number of zeros in ω. Since $f_\lambda(z)$ has only one zero in ω, we conclude that $z_n(\lambda)$ belongs to ω, and so the lemma is proved. $\square$

Proof of Lemma 6. We have from (3.36)

$$|z_n'(\lambda)|^2 = \frac{1}{a_n^2(\lambda) + b_n^2(\lambda)},$$

where a_n, b_n are defined in (3.37). Thus, using (3.39), we obtain

$$|z_n'(\lambda)| \le \frac{1}{|a_n(\lambda)|} \le \frac{(x_n^2(\lambda) + y_n^2(\lambda))^2}{2y_n^2(\lambda)(c_{m,n} - 1)}.$$

This and (3.62) prove that $|x_n'(\lambda)| < C$ for $\lambda \in U_\delta(\lambda_0)$. The inequality $|x_n'(\lambda)| > C$ was proved in Lemma 7. $\square$

References

[1] *L. Pastur, M. Shcherbina*, Universality of the local eigenvalue statistics for a class of unitary invariant random matrix ensembles. - J. Stat. Phys.(1997), 86, p.109-147

[2] *P. Deift, T. Kriecherbauer, K. McLaughlin, S. Venakides, X. Zhou*, Uniform asymptotics for polynomials orthogonal with respect to varying exponential weights and applications to universality questions in random matrix theory. - Commun. Pure Appl. Math.(1999), 52, p. 1335-1425

[3] *L. Pastur, M. Shcherbina*, Bulk Universality and related properties of Hermitian matrix model.- J.Stat.Phys.(2007), 130, p.205-250

[4] T. Tao, V. Vu "Random matrices: universality of local eigenvalue statistics", arXiv:0906.0510v6 [math.PR]

[5] L. Erdos, J. Ramirez, B. Schlein, H.-T. Yau, "Bulk universality for Wigner hermitian matrices", arXiv:0905.4176v1 [math-ph]

[6] L. Erdos, J. Ramirez, B. Schlein, T. Tao, V. Vu, H.-T. Yau, "Bulk universality for Wigner hermitian matrices with subexponential decay", arXiv:0906.4400v1 [math.PR]

[7] *V.A. Marchenko, L.A. Pastur,*, Distribution of eigenvalues for some sets of random matrices. - Math.USSR-Sb.(1967), 1, p.457-483.

[8] Nagao, T.; Wadati, M. Correlation functions of random matrix ensembles related to classical orthogonal polynomials. J. Phys. Soc. Japan 61 (1992), 1910Ц1918.

[9] *J.W.Silverstein, S.-I.Choi*, Analysis of the limiting spectral distribution of large dimensional random matrices. - J. Mult.Anal.(1995), 54, p.295-309

[10] *T.Shcherbina*, On Universality of Bulk Local Regime of the Deformed Gaussian Unitary Ensemble. -preprint

[11] *M.L.Mehta*, Random Matrices. -Academic Press, New York (1991)

[12] *E.Brezin, S. Hikami*, Correlation of nearby levels induced by a random potential.- Nucl.Phys.(1996),479, p.697-706

[13] *E.Brezin, S. Hikami*, Extension of level-spacing universality.- Phys.Rev. E(1997),56, p.264-269

[14] *E.Brezin, S. Hikami*, Level spacing of random matrices in an external source. -Phys.Rev. E (1998), 58, p. 7176-7185

[15] J.Baik, G.Ben Arours, S.Peche, Phase transition of the largest eigenvalue for nonnull complex sample covariance matrices".- Anal. of Prob.(2005), 33,5, p.1643-1697

[16] G.Ben Arours, S.Peche, Universality of local eigenvalue statistics for some sample covariance matrices.- Com. on P. and Apl.Math (2005), LVIII, p.1316-1357

[17] *K. Johansson*, Universality of the local spacing distribution in certain ensembles of Hermitian Wigner Matrices. -Commun. Math. Phys.(2001), 215, p.683-705

[18] *P.M.Bleher., A.B.J. Kuijlaars*,Large n limit of Gaussian random matrices with external source, part II, - Commun. Math.Phys. (2004),252, p.43-76

[19] *A.I. Aptekarev, P.M. Bleher, A.B.J. Kuijlaars*, Large n limit of Gaussian random matrices with external source, part II, -Commun. Math.Phys.(2005),25,p. 367-389

[20] R. Courant, D. Hilbert, *Methods of Mathematical Physics*, Vol. I, Interscience, NY, 1953.

[21] G. Polya, G. Szego, *Problems and theorems in analysis.* Die Grundlehren der math.Wissenschaften, Springer-Verlag, Vol. II, 1976